

SUPER EAT LABELINGS OF SUBDIVIDED STARS

M. JAVAID¹ AND A.A. BHATTI²

^{1,2}Department of Mathematics
National University of Computer and
Emerging Sciences Lahore, Pakistan.
javaidmath@gmail.com
akhlaq.ahmad@nu.edu.pk

Abstract. Kotzig and Rosa (1970) conjectured that every tree admits edge-magic total labeling. Enomoto et al. (1998) proposed the conjecture that every tree is super edge-magic total. In this paper, we describe super (a, d) -edge-antimagic total labelings on a subclass of the subdivided stars denoted by $T(n, n, n, n, n_5, n_6, \dots, n_r)$ for $d \in \{0, 1, 2\}$, where $n \geq 3$ odd, $r \geq 5$ and $n_m = 2^{m-4}(n-1) + 1$ for $5 \leq m \leq r$.

Key words: Super (a, d) -EAT labelings, subdivision of stars.

Abstrak. Kotzig dan Rosa (1970) telah membuat konjektur bahwa setiap *tree* dapat menghasilkan *edge-magic total labeling*. Enomoto et al. (1998) telah membuat konjektur bahwa setiap *tree* adalah *super edge-magic total*. Di dalam makalah ini, kami menjelaskan *super (a, d)-edge-antimagic total labeling* pada sebuah sub-kelas dari *star* yang terbagi yang dinyatakan oleh $T(n, n, n, n, n_5, n_6, \dots, n_r)$ untuk $d \in \{0, 1, 2\}$, dimana $n \geq 3$ ganjil, $r \geq 5$ dan $n_m = 2^{m-4}(n-1) + 1$ untuk $5 \leq m \leq r$.

Kata kunci: Super (a, d) -EAT labelings, pembagian stars.

1. INTRODUCTION

All graphs in this paper are finite, undirected and simple. For a graph G , $V(G)$ and $E(G)$ denote the vertex-set and the edge-set, respectively. A (v, e) -graph G is a graph such that $|V(G)| = v$ and $|E(G)| = e$. A general reference for graph-theoretic ideas can be seen in [29]. A *labeling* (or *valuation*) of a graph is a map that carries graph elements to numbers (usually to positive or non-negative integers). In this paper, the domain will be the set of all vertices and edges and such a labeling is called a *total labeling*. Some labelings use the vertex-set only or the

2000 Mathematics Subject Classification: 05C78 37M10.

Received: 17-11-2012, revised: 27-05-2013, accepted: 06-06-2013.

edge-set only and we shall call them *vertex-labelings* or *edge-labelings*, respectively. A number of classification studies on edge antimagic total graphs has been intensively investigated. For further studies on antimagic labelings, reader can see [13, 5].

Definition 1.1 A (s, d) -edge-antimagic vertex $((s, d)$ -EAV) labeling of a graph G is a bijective function $\lambda : V(G) \rightarrow \{1, 2, \dots, v\}$ such that the set of edge-sums of all edges in G , $\{w(xy) = \lambda(x) + \lambda(y) : xy \in E(G)\}$, forms an arithmetic progression $\{s, s + d, s + 2d, \dots, s + (e - 1)d\}$, where $s > 0$ and $d \geq 0$ are two fixed integers.

Definition 1.2. An (a, d) -edge-antimagic total $((a, d)$ -EAT) labeling of a graph G is a bijective function $\lambda : V(G) \cup E(G) \rightarrow \{1, 2, \dots, v + e\}$ such that the set of edge-weights of all edges in G , $\{w(xy) = \lambda(x) + \lambda(xy) + \lambda(y) : xy \in E(G)\}$, forms an arithmetic progression $\{a, a + d, a + 2d, \dots, a + (e - 1)d\}$, where $a > 0$ and $d \geq 0$ are two fixed integers. If such a labeling exists then G is said to be an (a, d) -EAT graph. Additionally, if $\lambda(V(G)) = \{1, 2, \dots, v\}$ then λ is called a super (a, d) -edge-antimagic total (super (a, d) -EAT) labeling and G becomes a super (a, d) -EAT graph.

In the above definition, if $d = 0$ then $(a, 0)$ -EAT labeling is called edge-magic total (EMT) labeling and super $(a, 0)$ -EAT labeling is called super edge-magic total (SEMT) labeling. The subject of edge-magic total (EMT) labeling of graphs has its origin in the works of Kotzig and Rosa [20, 21] on what they called magic valuations of graphs. The definition of (a, d) -EAT labeling was introduced by Simanjuntak, Bertault and Miller in [27] as a natural extension of EMT labeling defined by Kotzig and Rosa. A super (a, d) -EAT labeling is a natural extension of the notion of SEMT labeling defined by Enomoto, Lladó, Nakamigawa and Ringel in [9]. Moreover, they proposed the following conjecture:

Conjecture 1.1 *Every tree admits SEMT labeling [9].*

In the favour of this conjecture, many authors have proved the existence of SEMT labelings for various particular classes of trees for examples [1-8, 10-12, 14-17, 24, 25, 28, 29]. Lee and Shah [22] verified this conjecture by a computer search for trees with at most 17 vertices. However, this conjecture is still open. Bača et al. investigated the following relationship between (s, d) -EAV labeling and (a, d) -EAT labeling [3]:

Proposition 1.1. If a (v, e) -graph G has a (s, d) -EAV labeling then G admits

- (i) a super $(s + v + 1, d + 1)$ -EAT labeling,
- (ii) a super $(s + v + e, d - 1)$ -EAT labeling. ■

The notion of dual labeling has been introduced by Wallis [30]. The next lemma follows from the principal of duality, which is first studied by Baskoro [8].

Lemma 1.1 If g is a super edge-magic total labeling of G with the magic constant c , then the function $g_1 : V(G) \cup E(G) \rightarrow \{1, 2, \dots, v + e\}$ defined by

$$g_1(x) = \begin{cases} v + 1 - g(x), & \text{for } x \in V(G), \\ 2v + e + 1 - g(x), & \text{for } x \in E(G), \end{cases}$$

is also a super-magic total labeling of G with the magic constant $c_1 = 4v + e + 3 - c$. ■

Definition 1.3 For $n_i \geq 1$ and $r \geq 2$, let $G \cong T(n_1, n_2, \dots, n_r)$ be a graph obtained by inserting $n_i - 1$ vertices to each of the i^{th} edge of the star $K_{1,r}$, where $1 \leq i \leq r$. Thus, the graph $T(\underbrace{1, 1, \dots, 1}_{r\text{-times}})$ is a star $K_{1,r}$.

Subdivided stars form a particular class of trees and many authors have proved the antimagicness for various subclasses of subdivided stars as follows:

- Lu [23, 24] has called the subdivided star $T(m, n, k)$ as a three-path tree. Moreover, he has proved that it is a SEMT graph if n and m are odd with $k = n + 1$ or $k = n + 2$.
- Ngurah et al. [25] have proved that $T(m, n, k)$ is a SEMT graph if n and m are odd with $k = n + 3$ or $k = n + 4$.
- In [26], Salman et al. have found the results related to SEMT labelings on the subdivision of stars S_n^m for $m = 1, 2$, where $S_n^1 \cong T(\underbrace{2, 2, \dots, 2}_{n\text{-times}})$ and

$$S_n^2 \cong T(\underbrace{3, 3, \dots, 3}_{n\text{-times}}).$$

- In [16], Javaid et al. have formulated SEMT labelings on the subdivision of star $K_{1,4}$ and w-trees.
- Javaid and Akhlaq [17] have proved that the subdivided stars $T(n, n, n + 2, n + 2, n_5, \dots, n_p)$ admit super (a, d) -EAT labelings, where $n \geq 3$ is odd, $r \geq 5$ and $n_m = 1 + (n + 1)2^{m-4}$ for $5 \leq m \leq r$.

However, the problem to find super (a, d) -EAT labelings on $T(n_1, n_2, n_3, \dots, n_r)$ for different $\{n_i : 1 \leq i \leq r\}$ is still open. In this paper, for $d \in \{0, 1, 2\}$, we find super (a, d) -EAT labelings on the subdivided stars $T(n, n, n, n, n_5, n_6, \dots, n_r)$, where $n \geq 3$ is odd, $r \geq 5$ and $n_m = 2^{m-4}(n - 1) + 1$ for $5 \leq m \leq r$.

2. BOUNDS OF MAGIC CONSTANT

In this section, we present different lemmas related to lower and upper bounds of the magic constant a for various subclasses of subdivided stars.

Ngurah et al. [25] found the following lower and upper bounds of the magic constant a for a particular subclass of the subdivided stars denoted by $T(m, n, k)$, which is given below:

Lemma 2.1. If $T(m, n, k)$ is a super $(a, 0)$ -EAT graph, then $\frac{1}{2l}(5l^2 + 3l + 6) \leq a \leq \frac{1}{2l}(5l^2 + 11l - 6)$, where $l = m + n + k$.

The lower and upper bounds of the magic constant a for a particular subclass of the subdivided stars $T(\underbrace{n, n, \dots, n}_{n\text{-times}})$ are established by Salman et al. [26] as follows:

Lemma 2.2. If $T(\underbrace{n, n, \dots, n}_{n\text{-times}})$ is a super $(a, 0)$ -EAT graph, then $\frac{1}{2l}(5l^2 + (9 - 2n)l + n^2 - n) \leq a \leq \frac{1}{2l}(5l^2 + (2n + 5)l + n - n^2)$, where $l = n^2$. ■

Javid [19] has proved lower and upper bounds of the magic constant a for the most extended subclasses of the subdivided stars denoted by $T(n_1, n_2, n_3, \dots, n_r)$ with any $n_i \geq 1$ for $1 \leq i \leq r$, which is presented in the following lemma:

Lemma 2.3. If $T(n_1, n_2, n_3, \dots, n_r)$ is a super $(a, 0)$ -EAT graph, then $\frac{1}{2l}(5l^2 + r^2 - 2lr + 9l - r) \leq a \leq \frac{1}{2l}(5l^2 - r^2 + 2lr + 5l + r)$, where $l = \sum_{i=1}^r n_i$. ■

3. SUPER (a, d) -EAT LABELINGS OF SUBDIVIDED STARS

In this section, we prove the main results related to super (a, d) -EAT labelings on a particular subclass of the subdivided stars for different values of the parameter d .

Theorem 2.1. For any odd $n \geq 3$, $G \cong T(n, n, n, n, 2n - 1)$ admits a super $(a, 0)$ -EAT labeling with $a = 2v + s - 1$ and a super $(\hat{a}, 2)$ -EAT labeling with $\hat{a} = v + s + 1$, where $v = |V(G)|$ and $s = 3n + 4$.

PROOF. Let us denote the vertices and edges of G , as follows:

$V(G) = \{c\} \cup \{x_i^{l_i} \mid 1 \leq i \leq 5; 1 \leq l_i \leq n_i\}$, $E(G) = \{c x_i^1 \mid 1 \leq i \leq 5\} \cup \{x_i^{l_i} x_i^{l_i+1} \mid 1 \leq i \leq 5; 1 \leq l_i \leq n_i - 1\}$. If $v = |V(G)|$ and $e = |E(G)|$ then $v = 6n$, and $e = 6n - 1$. Now, we define the labeling $\lambda : V(G) \rightarrow \{1, 2, \dots, v\}$ as follows:

$$\lambda(c) = 4n + 2.$$

For $1 \leq l_i \leq n_i$ odd;

$$\lambda(u) = \begin{cases} \frac{l_1+1}{2}, & \text{for } u = x_1^{l_1}, \\ (n+2) - \frac{l_2+1}{2}, & \text{for } u = x_2^{l_2}, \\ (n+1) + \frac{l_3+1}{2}, & \text{for } u = x_3^{l_3}, \\ (2n+3) - \frac{l_4+1}{2}, & \text{for } u = x_4^{l_4}, \\ (3n+3) - \frac{l_5+1}{2}, & \text{for } u = x_5^{l_5}. \end{cases}$$

For $2 \leq l_i \leq n_i - 1$ even;

$$\lambda(u) = \begin{cases} (3n+2) + \frac{l_1}{2}, & \text{for } u = x_1^{l_1}, \\ (4n+2) - \frac{l_2}{2}, & \text{for } u = x_2^{l_2}, \\ (4n+2) + \frac{l_3}{2}, & \text{for } u = x_3^{l_3}, \\ (5n+2) - \frac{l_4}{2}, & \text{for } u = x_4^{l_4}, \\ (6n+1) - \frac{l_5}{2}, & \text{for } u = x_5^{l_5}. \end{cases}$$

The set of all edge-sums generated by the above formulas forms a consecutive integer sequence $s = 3n+4, 3n+5, \dots, 3n+3+e$. Therefore, by Proposition 1.1, λ can be extended to a super $(a, 0)$ -EAT labeling with magic constant $a = 2v+s-1 = 15n+3$ and to a super $(\acute{a}, 2)$ -EAT labeling with minimum edge-weight $\acute{a} = v+1+s = 9n+5$. ■

Theorem 2.2. For any odd $n \geq 3$, $G \cong T(n, n, n, n, 2n-1)$ admits a super $(a, 1)$ -EAT labeling with $a = s + \frac{3v}{2}$, where $v = |V(G)|$ and $s = 3n+4$.

PROOF. Let us consider the vertex and edge set of G and the labeling $\lambda : V(G) \rightarrow \{1, 2, \dots, v\}$ by the same manner as in Theorem 2.1. It follows that edge-sums of all the edges of G constitute an arithmetic sequence $3n+4, 3n+5, \dots, 3n+3+e$, with common difference 1. We denote it by $A = \{a_i; 1 \leq i \leq e\}$. Now to show that λ is an $(a, 1)$ -EAT labeling of G , define the set of edge-labels as $B = \{b_j = v + j; 1 \leq j \leq e\}$. The set of edge-weights can be obtained as $C = \{a_{2i-1} + b_{e-i+1}; 1 \leq i \leq \frac{e+1}{2}\} \cup \{a_{2j} + b_{\frac{e-1}{2}-j+1}; 1 \leq j \leq \frac{e-1}{2}\}$. It is easy to see that C constitutes an arithmetic sequence with $d = 1$ and $a = s + \frac{3v}{2} = 12n+4$. Since all vertices receive the smallest labels, λ is a super $(a, 1)$ -EAT labeling. ■

Theorem 2.3. For any odd $n \geq 3$, $G \cong T(n, n, n, n, 2n-1, 4n-3)$ admits a super $(a, 0)$ -EAT labeling with $a = 2v+s-1$ and a super $(\acute{a}, 2)$ -EAT labeling with $\acute{a} = v+s+1$, where $v = |V(G)|$ and $s = 5n+3$.

PROOF. Let us denote the vertices and edges of G , as follows:

$V(G) = \{c\} \cup \{x_i^{l_i} \mid 1 \leq i \leq 6; 1 \leq l_i \leq n_i\}$, $E(G) = \{c x_i^1 \mid 1 \leq i \leq 6\} \cup \{x_i^{l_i} x_i^{l_i+1} \mid 1 \leq i \leq 6; 1 \leq l_i \leq n_i - 1\}$. If $v = |V(G)|$ and $e = |E(G)|$ then $v = 10n - 3$, and $e = 10n - 4$. Now, we define the labeling $\lambda : V(G) \rightarrow \{1, 2, \dots, v\}$ as follows:

$$\lambda(c) = 6n + 1.$$

For $1 \leq l_i \leq n_i$ odd;

$$\lambda(u) = \begin{cases} \frac{l_1+1}{2}, & \text{for } u = x_1^{l_1}, \\ (n+2) - \frac{l_2+1}{2}, & \text{for } u = x_2^{l_2}, \\ (n+1) + \frac{l_3+1}{2}, & \text{for } u = x_3^{l_3}, \\ (2n+3) - \frac{l_4+1}{2}, & \text{for } u = x_4^{l_4}, \\ (3n+3) - \frac{l_5+1}{2}, & \text{for } u = x_5^{l_5}, \\ (5n+2) - \frac{l_6+1}{2}, & \text{for } u = x_6^{l_6}. \end{cases}$$

For $2 \leq l_i \leq n_i - 1$ even;

$$\lambda(u) = \begin{cases} (5n+1) + \frac{l_1}{2}, & \text{for } u = x_1^{l_1}, \\ (6n+1) - \frac{l_2}{2}, & \text{for } u = x_2^{l_2}, \\ (6n+1) + \frac{l_3}{2}, & \text{for } u = x_3^{l_3}, \\ (7n+1) - \frac{l_4}{2}, & \text{for } u = x_4^{l_4}, \\ 8n - \frac{l_5}{2}, & \text{for } u = x_5^{l_5}, \\ (10n-2) - \frac{l_6}{2}, & \text{for } u = x_6^{l_6}. \end{cases}$$

The set of all edge-sums generated by the above formulas forms a consecutive integer sequence $s = 5n + 3, 5n + 4, \dots, 5n + 2 + e$. Therefore, by Proposition 1.1, λ can be extended to a super $(a, 0)$ -EAT labeling with magic constant $a = 2v + s - 1 = 25n - 4$ and to a super $(\acute{a}, 2)$ -EAT labeling with minimum edge-weight $\acute{a} = v + 1 + s = 15n + 1$. \blacksquare

Theorem 2.4. For any odd $n \geq 3$, $G \cong T(n, n, n, n, 2n - 1, 4n - 3, 8n - 7)$ admits a super $(a, 0)$ -EAT labeling with $a = 2v + s - 1$ and a super $(\acute{a}, 2)$ -EAT labeling with $\acute{a} = v + s + 1$, where $v = |V(G)|$ and $s = 9n$.

PROOF. Let us denote the vertices and edges of G , as follows:

$V(G) = \{c\} \cup \{x_i^{l_i} \mid 1 \leq i \leq 7; 1 \leq l_i \leq n_i\}$, $E(G) = \{c x_i^1 \mid 1 \leq i \leq 7\} \cup \{x_i^{l_i} x_i^{l_i+1} \mid 1 \leq i \leq 7; 1 \leq l_i \leq n_i - 1\}$. If $v = |V(G)|$ and $e = |E(G)|$ then $v =$

$18n - 10$, and $e = 18n - 11$. Now, we define the labeling $\lambda : V(G) \rightarrow \{1, 2, \dots, v\}$ as follows:

$$\lambda(c) = 10n - 2.$$

For $1 \leq l_i \leq n_i$ odd;

$$\lambda(u) = \begin{cases} \frac{l_1+1}{2}, & \text{for } u = x_1^{l_1}, \\ (n+2) - \frac{l_2+1}{2}, & \text{for } u = x_2^{l_2}, \\ (n+1) + \frac{l_3+1}{2}, & \text{for } u = x_3^{l_3}, \\ (2n+3) - \frac{l_4+1}{2}, & \text{for } u = x_4^{l_4}, \\ (3n+3) - \frac{l_5+1}{2}, & \text{for } u = x_5^{l_5}, \\ (5n+2) - \frac{l_6+1}{2}, & \text{for } u = x_6^{l_6}, \\ (9n-1) - \frac{l_7+1}{2}, & \text{for } u = x_7^{l_7}. \end{cases}$$

For $2 \leq l_i \leq n_i - 1$ even;

$$\lambda(u) = \begin{cases} (9n-2) + \frac{l_1}{2}, & \text{for } u = x_1^{l_1}, \\ (10n-2) - \frac{l_2}{2}, & \text{for } u = x_2^{l_2}, \\ (10n-2) + \frac{l_3}{2}, & \text{for } u = x_3^{l_3}, \\ (11n-2) - \frac{l_4}{2}, & \text{for } u = x_4^{l_4}, \\ (12n-3) - \frac{l_5}{2}, & \text{for } u = x_5^{l_5}, \\ (14n-5) - \frac{l_6}{2}, & \text{for } u = x_6^{l_6}, \\ (18n-9) - \frac{l_7}{2}, & \text{for } u = x_7^{l_7}. \end{cases}$$

The set of all edge-sums generated by the above formulas forms a consecutive integer sequence $s = 9n, 9n+1, \dots, 9n-1+e$. Therefore, by Proposition 1.1, λ can be extended to a super $(a, 0)$ -EAT labeling with magic constant $a = 2v+s-1 = 45n-21$ and to a super $(\acute{a}, 2)$ -EAT labeling with minimum edge-weight $\acute{a} = v+1+s = 27n-9$. ■

Theorem 2.5. For any odd $n \geq 3$, $G \cong T(n, n, n, n, 2n-1, 4n-3, 8n-7)$ admits a super $(a, 1)$ -EAT labeling with $a = s + \frac{3v}{2}$, where $v = |V(G)|$ and $s = 9n$.

PROOF. Let us consider the vertex and edge set of G and the labeling $\lambda : V(G) \rightarrow \{1, 2, \dots, v\}$ by the same manner as in Theorem 2.4. It follows that edge-sums of all the edges of G constitute an arithmetic sequence $9n, 9n+1, \dots, 9n-1+e$, with

common difference 1. We denote it by $A = \{a_i; 1 \leq i \leq e\}$. Now to show that λ is an $(a, 1)$ -EAT labeling of G , define the set of edge-labels as $B = \{b_j = v + j; 1 \leq j \leq e\}$. The set of edge-weights can be obtained as $C = \{a_{2i-1} + b_{e-i+1}; 1 \leq i \leq \frac{e+1}{2}\} \cup \{a_{2j} + b_{\frac{e-1}{2}-j+1}; 1 \leq j \leq \frac{e+1}{2} - 1\}$. It is easy to see that C constitutes an arithmetic sequence with $d = 1$ and $a = s + \frac{3v}{2} = 36n - 15$. Since all vertices receive the smallest labels, λ is a super $(a, 1)$ -EAT labeling. ■

Theorem 2.6. For any $n \geq 3$ odd, $G \cong T(n, n, n, n, n_5, \dots, n_r)$ admits a super $(a, 0)$ -EAT labeling with $a = 2v + s - 1$ and a super $(\acute{a}, 2)$ -EAT labeling with $\acute{a} = v + s + 1$ where $v = |V(G)|$, $s = (2n + 4) + \sum_{m=5}^r [2^{m-5}(n-1) + 1]$, $r \geq 5$ and $n_m = 2^{m-4}(n-1) + 1$ for $5 \leq m \leq r$.

PROOF. Let us denote the vertices and edges of G , as follows:

$V(G) = \{c\} \cup \{x_i^{l_i} \mid 1 \leq i \leq r; 1 \leq l_i \leq n_i\}$, $E(G) = \{c x_i^1 \mid 1 \leq i \leq r\} \cup \{x_i^{l_i} x_i^{l_i+1} \mid 1 \leq i \leq r; 1 \leq l_i \leq n_i - 1\}$. If $v = |V(G)|$ and $e = |E(G)|$ then $v = (4n + 1) + \sum_{m=5}^r [2^{m-4}(n-1) + 1]$ and $e = v - 1$. Now, we define the labeling $\lambda : V(G) \rightarrow \{1, 2, \dots, v\}$ as follows:

$$\lambda(c) = (3n + 2) + \sum_{m=5}^r [2^{m-5}(n-1) + 1].$$

For $1 \leq l_i \leq n_i$ odd, where $i = 1, 2, 3, 4$ and $5 \leq i \leq r$, we define

$$\lambda(u) = \begin{cases} \frac{l_1+1}{2}, & \text{for } u = x_1^{l_1}, \\ (n+2) - \frac{l_2+1}{2}, & \text{for } u = x_2^{l_2}, \\ (n+1) + \frac{l_3+1}{2}, & \text{for } u = x_3^{l_3}, \\ (2n+3) - \frac{l_4+1}{2}, & \text{for } u = x_4^{l_4}. \end{cases}$$

$$\lambda(x_i^{l_i}) = (2n + 3) + \sum_{m=5}^i [2^{m-5}(n-1) + 1] - \frac{l_i + 1}{2} \text{ respectively.}$$

Let $\alpha = (2n + 2) + \sum_{m=5}^r [2^{m-5}(n-1) + 1]$. For $2 \leq l_i \leq n_i$ even, and $1 \leq i \leq r$, we define

$$\lambda(u) = \begin{cases} \alpha + \frac{l_1}{2}, & \text{for } u = x_1^{l_1}, \\ (\alpha + n) - \frac{l_2}{2}, & \text{for } u = x_2^{l_2}, \\ (\alpha + n) + \frac{l_3}{2}, & \text{for } u = x_3^{l_3}, \\ (\alpha + 2n) - \frac{l_4}{2}, & \text{for } u = x_4^{l_4}. \end{cases}$$

and

$$\lambda(x_i^{l_i}) = (\alpha + 2n) + \sum_{m=5}^i [2^{m-5}(n-1)] - \frac{l_i}{2}.$$

The set of all edge-sums generated by the above formulas forms a consecutive integer sequence $s = \alpha + 2, \alpha + 3, \dots, \alpha + 1 + e$. Therefore, by Proposition 1.1, λ can be extended to a super $(a, 0)$ -EAT labeling with magic constant $a = v + e + s = 2v + (2n + 3) + \sum_{m=5}^r [2^{m-5}(n-1) + 1]$ and to a super $(\acute{a}, 2)$ -EAT labeling with minimum edge-weight $\acute{a} = v + 1 + s = v + (2n + 5) + \sum_{m=5}^r [2^{m-5}(n-1) + 1]$. ■

THEOREM 2.7. For any $n \geq 3$ odd, $G \cong T(n, n, n, n, n_5, \dots, n_r)$ admits super $(a, 1)$ -EAT labeling with $a = s + \frac{3v}{2}$ if v is even, where $v = |V(G)|$, $s = (2n + 4) + \sum_{m=5}^r [2^{m-5}(n-1) + 1]$, $r \geq 5$, and $n_m = 2^{m-4}(n-1) + 1$ for $5 \leq m \leq r$.

PROOF. Let us consider the vertex and edge set of G and the labeling $\lambda : V(G) \rightarrow \{1, 2, \dots, v\}$ by the same manner as in Theorem 2.6. It follows that edge-sums of all the edges of G constitute an arithmetic sequence $s = \alpha + 2, \alpha + 3, \dots, \alpha + 1 + e$ with common difference 1, where $\alpha = (2n + 2) + \sum_{m=5}^r [2^{m-5}(n-1) + 1]$. We denote it by $A = \{a_i; 1 \leq i \leq e\}$. Now to show that λ is an $(a, 1)$ -EAT labeling of G , define the set of edge-labels as $B = \{b_j = v + j; 1 \leq j \leq e\}$. The set of edge-weights can be obtained as $C = \{a_{2i-1} + b_{e-i+1}; 1 \leq i \leq \frac{e+1}{2}\} \cup \{a_{2j} + b_{\frac{e-1}{2}-j+1}; 1 \leq j \leq \frac{e+1}{2} - 1\}$. It is easy to see that C constitutes an arithmetic sequence with $d = 1$ and $a = s + \frac{3v}{2}$. Since, all vertices receive the smallest labels, λ is a super $(a, 1)$ -EAT labeling. ■

From Theorems 2.1, 2.3, 2.4 and 2.6 by the principal of duality it follows that we can find the super $(a, 0)$ -EAT labelings with different magic constant. Thus, we have the following corollaries:

Corollary 2.1. For any odd $n \geq 3$, $T(n, n, n, n, 2n - 1)$ admits a super $(a, 0)$ -EAT total labeling with magic constant $a = 15n - 1$.

Corollary 2.2. For any odd $n \geq 3$, $T(n, n, n, n, 2n - 1, 4n - 3)$ admits a super $(a, 0)$ -EAT labeling with magic constant $a = 25n - 9$.

Corollary 2.3. For any odd $n \geq 3$, $T(n, n, n, n, 2n - 1, 4n - 3, 8n - 7)$ admits a super $(a, 0)$ -EAT labeling with magic constant $a = 45n - 27$.

Corollary 2.4. For any $n \geq 3$ odd, and $r \geq 5$, $T(n, n, n, n, n_5, \dots, n_r)$ admits a super $(a, 0)$ -EAT total labeling with $a = 3v - (2n + 1) - \sum_{m=5}^r [2^{m-5}(n - 1) + 1]$, where $n_m = 2^{m-4}(n - 1) + 1$ for $5 \leq m \leq r$.

4. CONCLUSION

In this paper, we have proved that a subclass of subdivided stars denoted by $T(n, n, n, n, n_5, \dots, n_r)$, admits super (a, d) -EAT labelings for $d = 0, 1, 2$, when $n \geq 3$ is odd, $r \geq 5$ and $n_m = 2^{m-4}(n - 1) + 1$ for $5 \leq m \leq r$.

Acknowledgement. The research contents of this paper are partially supported by the Higher Education Commission (HEC) of Pakistan and National University of Computer and Emerging Sciences (NUCES) Lahore, Pakistan.

References

- [1] Bača, M., Lin, Y., and Muntaner-Batle, F.A., "Edge-antimagic labeling of forests", *Utilitas Math.*, **81** (2010), 31-40.
- [2] Bača, M., Lin, Y., Miller, M., and Youssef, M.Z., "Edge-antimagic graphs", *Discrete Math.*, **307** (2007), 1232-1244.
- [3] Bača, M., Lin, Y., Miller, M., and Simanjuntak, R., "New constructions of magic and antimagic graph labelings", *Utilitas Math.*, **60** (2001), 229-239.
- [4] Bača, M., Lin, Y., and Muntaner-Batle, F.A., "Super edge-antimagic labelings of the path-like trees", *Utilitas Math.*, **73** (2007), 117-128.
- [5] Bača, M., and Miller, M., *Super Edge-Antimagic Graphs*, Brown Walker Press, Boca Raton, Florida USA, 2008.
- [6] Bača, M., Semaničová -Feňovčíková, A., and Shafiq, M.K., "A method to generate large classes of edge-antimagic trees", *Utilitas Math.*, **86** (2011), 33-43.
- [7] Baskoro, E.T., and Ngurah, A.A.G., "On super edge-magic total labelings", *Bull. Inst. Combin. Appl.*, **37** (2003), 82-87.
- [8] Baskoro, E.T., Sudarsana, I.W., and Cholily, Y.M., "How to construct new super edge-magic graphs from some old ones", *J. Indones. Math. Soc. (MIHMI)*, **11:2** (2005), 155-162.
- [9] Enomoto, H., Llado, A.S., Nakamigawa, T., and Ringel, G., "Super edge-magic graphs", *SUT J. Math.*, **34** (1980), 105-109.
- [10] Figueroa-Centeno, R.M., Ichishima, R., and Muntaner-Batle, F.A., "The place of super edge-magic labeling among other classes of labeling", *Discrete Math.*, **231** (2001), 153-168.
- [11] Figueroa-Centeno, R.M., Ichishima, R., and Muntaner-Batle, F.A., "On super edge-magic graph", *Ars Combin.*, **64** (2002), 81-95.
- [12] Fukuchi, Y., "A recursive theorem for super edge-magic labeling of trees", *SUT J. Math.*, **36** (2000), 279-285.
- [13] Gallian, J.A., "A dynamic survey of graph labeling", *Electron. J. Combin.*, (2012).
- [14] Hussain, M., Baskoro, E.T., and Slamim, "On super edge-magic total labeling of banana trees", *Utilitas Math.*, **79** (2009), 243-251.

- [15] Javaid, M., Bhatti, A.A., Hussain, M., "On (a, d) -edge-antimagic total labeling of extended w-trees", *Utilitas Math.*, **87** (2012), 293-303.
- [16] Javaid, M., Hussain, M., Ali, K., and Shaker, H., "Super edge-magic total labeling on subdivision of trees", *Utilitas Math.*, **89** (2012), 169 - 177.
- [17] Javaid, M., and Bhatti, A.A., On super (a, d) -edge-antimagic total labeling of subdivided stars. *Ars Combin.*, **105** (2012), 503-512.
- [18] Javaid, M., Hussain, M., Ali, K., and Dar, K.H., "Super edge-magic total labeling on w-trees", *Utilitas Math.*, **86** (2011), 183-191.
- [19] Javaid, M., *Labeling Graphs and Hypergraphs*, PhD Thesis FAST-NUCES, Lahore Campus, Pakistan, 2013.
- [20] Kotzig, A., and Rosa, A., Magic valuations of finite graphs, *Canad. Math. Bull.*, **13** (1970), 451-461.
- [21] Kotzig, A., and Rosa, A., *Magic valuation of complete graphs*, Centre de Recherches Mathematiques, Universite de Montreal, 1972, CRM-175.
- [22] Lee, S.M., and Shah, Q.X., "All trees with at most 17 vertices are super edge-magic", *16th MCCC Conference, Carbondale*, University Southern Illinois, (2002).
- [23] Lu, Y.J., "A proof of three-path trees $P(m, n, t)$ being edge-magic", *College Mathematica*, **17:2** (2001), 41-44.
- [24] Lu, Y.J., "A proof of three-path trees $P(m, n, t)$ being edge-magic (II)", *College Mathematica*, **20:3** (2004), 51-53.
- [25] Ngurah, A.A.G., Simanjuntak, R., and Baskoro, E.T., "On (super) edge-magic total labeling of subdivision of $K_{1,3}$ ", *SUT J. Math.*, **43** (2007), 127-136.
- [26] Salman, A.N.M., Ngurah, A.A.G., and Izzati, N., On "Super Edge-Magic Total Labeling of a Subdivision of a Star S_n ", *Utilitas Mathematica*, **81** (2010), 275-284.
- [27] Simanjuntak, R., Bertault, F., and Miller, M., "Two new (a, d) -antimagic graph labelings", *Proc. of Eleventh Australasian Workshop on Combinatorial Algorithms*, (2000), 179-189.
- [28] Sugeng, K.A., Miller, M., Slamin, and Bača, M., " (a, d) -edge-antimagic total labelings of caterpillars", *Lecture Notes Comput. Sci.*, **3330** (2005), 169-180.
- [29] West, D.B., An Introduction to Graph Theory *Prentice-Hall*, 1996.
- [30] Wallis, W.D., Magic Graphs, *Birkhäuser*, Boston-Basel-Berlin, 2001.