

Existence of Fixed Points in Neutrosophic Bipolar Fuzzy Metric Spaces with Application

Vishal Gupta¹, Anju^{2*}, Deepti Thakur³, and Ishit Sehgal⁴

^{1,2,4}Department of Mathematics, Maharishi Markandeshwar (Deemed to be University),
India,

¹vishal.gmn@gmail.com, ²anju.kaliraman02@gmail.com, ⁴ishitsehgal06@gmail.com

³Mathematics Preparatory Studies Centre, University of Technology and Applied
Sciences, Oman, deepti.thakur@utas.edu.com].

Abstract. In this paper, we aim to introduce and explain the concept of neutrosophic fuzzy bipolar metric spaces and provide some fixed point results in this new setting. Additionally, we give a detailed presentation of the proof of fixed point theorems using covariant maps, showing how our findings extend and generalize those established in the existing body of literature. The derived results are supported with a suitable example and an application.

Key words and Phrases: Neutrosophic fuzzy bipolar metric space, fixed point, covariant mapping.

1. INTRODUCTION

Zadeh [1] was the pioneer behind the introduction of fuzzy sets. Fuzzy theory's rapid evolution has simplified its application, enabling us to explore uncertainty levels in nature through a rigorous and mathematical approach. The incorporation of both probabilistic metric space and fuzzy concepts led to the introduction of fuzzy metric space. This pioneering work, attributed to Kaleva and Seikkala [2] characterizes FMS as a representation of distance between two points through non-negative fuzzy numbers.

FMS has since found practical applications in diverse fields, including fixed point theory, image and signal processing, medical imaging, and decision-making. Schweizer and Sklar [3] introduced the concept of continuous t-norms, which laid the foundation for further developments in the field. Kramosil and Michalek [4] extended the study by comparing probabilistic and statistical generalizations of

*Corresponding author

2020 Mathematics Subject Classification: 47H10, 54H25.

Received: 19-03-2024, accepted: 09-07-2025.

metric spaces, incorporating the concept of fuzziness. Garbiec [5] interpreted the fuzzy concept of Banach contraction, and this idea has since been explored in various contexts.

With the emergence of intuitionistic fuzzy sets (IFS), these versatile sets have permeated the various domains of fuzzy set theory. Park [6] extends this reach by defining the intuitionistic Fuzzy Metric Space (IFMS), a broadened idia of FMS. Park [6] incorporates George and Veeramani's [7] insights by applying t-norm and t-conorm operations to FMS, effectively defining IFMS and exploring its foundational characteristics. Many researchers work on FMS and IFMS, see references [8, 9, 10]. In 2024 Gupta, Saini and Gondhi [11] combine MIF metric with soft and proved FP theorem in this newly defined space, also see [12, 13].

Mutlu and Gurdal [14] introduced bipolar metric spaces and proved fixed point theorems, initiating a trend in utilizing different types of contractions in bipolar metric spaces, as evidenced by various researchers, see in references [15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26]. The utilization of intuitionistic fuzzy bipolar metric spaces for solving integral equations is another noteworthy development, emphasizing membership functions within fuzzy metric spaces.

Kirişci and Simşek [27] introduced the concept of neutrosophic metric spaces (NMSs), which extended the framework of intuitionistic fuzzy metric spaces by incorporating the principles of neutrosophy. They explored various topological properties of these spaces and established numerous fixed point results for contraction mappings within NMSs. Several other researchers have also pursued related advancements, enriching the theory with additional generalizations and applications, see [28, 29, 30, 31].

The paper's structure is as follows: The first section provides an introduction, the second section covers basic definitions in the preliminaries, the third section presents the neutrosophic fuzzy bipolar metric space and some fixed-point results, accompanied by illustrative examples. In the fourth section, we demonstrate applications, including the proof of existence and uniqueness of a solution for a first-order ordinary differential equation.

2. PRELIMINARIES

Definition 2.1. [15] *A binary operation $*$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ is called a continuous triangle norm (CTN) if:*

- (1) $\varpi * \aleph = \aleph * \varpi$, for all $\varpi, \aleph \in [0, 1]$;
- (2) $*$ is continuous;
- (3) $\varpi * 1 = \varpi$, for all $\varpi \in [0, 1]$;
- (4) $(\varpi * \aleph) * \omega = \varpi * (\aleph * \omega)$, for all $\varpi, \aleph, \omega \in [0, 1]$;
- (5) If $\varpi \leq \aleph$ and $\zeta \leq \omega$, with $\varpi, \aleph, \zeta, \omega \in [0, 1]$, then $\varpi * \zeta \leq \omega * \aleph$.

Definition 2.2. [15] *A continuous triangle co-norm (CTCN) is defined as a binary operation $\circ$: $[0, 1] \times [0, 1] \rightarrow [0, 1]$ satisfying the following properties:*

- (1) $\varpi \circ \aleph = \aleph \circ \varpi$, for all $\varpi, \aleph \in [0, 1]$;
- (2) The operation $\circ$ is continuous;
- (3) $\varpi \circ 0 = 0$, for all $\varpi \in [0, 1]$;
- (4) $(\varpi \circ \aleph) \circ \omega = \varpi \circ (\aleph \circ \omega)$, for all $\varpi, \aleph, \omega \in [0, 1]$;
- (5) If $\varpi \leq \aleph$ and $\zeta \leq \omega$, with $\varpi, \aleph, \zeta, \omega \in [0, 1]$, then $\varpi \circ \zeta \leq \aleph \circ \omega$.

Definition 2.3. [4] Let $*$ be a CTN, $\circ$ be a CTCN, and $\mathbb{H}, \mathbb{O}$ be fuzzy sets on $\Omega \times \Omega \times (0, +\infty)$, where $\Omega \neq \emptyset$. If $(\Omega, \mathbb{H}, \mathbb{O}, *, \circ)$ fulfills the below conditions for all $\aleph, \varpi \in \Omega$ and $\psi, \varphi > 0$:

- (1) $\mathbb{H}(\aleph, \varpi, \varphi) + \mathbb{O}(\aleph, \varpi, \varphi) \leq 1$;
- (2) $\mathbb{H}(\aleph, \varpi, \varphi) > 0$;
- (3) $\mathbb{H}(\aleph, \varpi, \varphi) = 1$ if and only if $\aleph = \varpi$;
- (4) $\mathbb{H}(\aleph, \varpi, \varphi) = \mathbb{H}(\varpi, \aleph, \varphi)$;
- (5) $\mathbb{H}(\aleph, \aleph_0, (\varphi + \psi)) \geq \mathbb{H}(\aleph, \varpi, \varphi) * \mathbb{H}(\varpi, \aleph_0, \psi)$;
- (6) $\mathbb{H}(\aleph, \varpi, \cdot)$ is an increasing function of $\mathbf{R}^+$ and $\lim_{\varphi \rightarrow +\infty} \mathbb{H}(\aleph, \varpi, \varphi) = 1$;
- (7) $\mathbb{O}(\aleph, \varpi, \varphi) > 0$;
- (8) $\mathbb{O}(\aleph, \varpi, \varphi) = 0$ if and only if $\aleph = \varpi$;
- (9) $\mathbb{O}(\aleph, \varpi, \varphi) = \mathbb{O}(\varpi, \aleph, \varphi)$;
- (10) $\mathbb{O}(\aleph, \aleph_0, (\varphi + \psi)) \leq \mathbb{O}(\aleph, \varpi, \varphi) \circ \mathbb{O}(\varpi, \aleph_0, \psi)$;
- (11) $\mathbb{O}(\aleph, \varpi, \cdot)$ is a decreasing function of $\mathbf{R}^+$ and $\lim_{\varphi \rightarrow +\infty} \mathbb{O}(\aleph, \varpi, \varphi) = 0$.

Then, $(\Omega, \mathbb{H}, \mathbb{O}, *, \circ)$ is an intuitionistic fuzzy metric space.

Definition 2.4. [4] Let $*$ be a CTN, $\circ$ be a CTCN, $b \geq 1$ and $\mathbb{H}, \mathbb{O}$ be fuzzy sets on $\Omega \times \Omega \times (0, +\infty)$ where $\Omega \neq \emptyset$. If $(\Omega, \mathbb{H}, \mathbb{O}, *, \circ, b)$ fulfills the below conditions for all $\aleph, \varpi \in \Omega$ and $\psi, \varphi > 0$:

- (1) $\mathbb{H}(\aleph, \varpi, \varphi) + \mathbb{O}(\aleph, \varpi, \varphi) \leq 1$;
- (2) $\mathbb{H}(\aleph, \varpi, \varphi) > 0$;
- (3) $\mathbb{H}(\aleph, \varpi, \varphi) = 1$ if and only if $\aleph = \varpi$;
- (4) $\mathbb{H}(\aleph, \varpi, \varphi) = \mathbb{H}(\varpi, \aleph, \varphi)$;
- (5) $\mathbb{H}(\aleph, \aleph_0, b(\varphi + \psi)) \geq \mathbb{H}(\aleph, \varpi, \varphi) * \mathbb{H}(\varpi, \aleph_0, \psi)$;
- (6) $\mathbb{H}(\aleph, \varpi, \cdot)$ is an increasing function of $\mathbf{R}^+$ and $\lim_{\varphi \rightarrow +\infty} \mathbb{H}(\aleph, \varpi, \varphi) = 1$;
- (7) $\mathbb{O}(\aleph, \varpi, \varphi) > 0$;
- (8) $\mathbb{O}(\aleph, \varpi, \varphi) = 0$ if and only if $\aleph = \varpi$;
- (9) $\mathbb{O}(\aleph, \varpi, \varphi) = \mathbb{O}(\varpi, \aleph, \varphi)$;
- (10) $\mathbb{O}(\aleph, \aleph_0, b(\varphi + \psi)) \leq \mathbb{O}(\aleph, \varpi, \varphi) \circ \mathbb{O}(\varpi, \aleph_0, \psi)$;
- (11) $\mathbb{O}(\aleph, \varpi, \cdot)$ is a decreasing function of $\mathbf{R}^+$ and $\lim_{\varphi \rightarrow +\infty} \mathbb{O}(\aleph, \varpi, \varphi) = 0$.

Then, $(\Omega, \mathbb{H}, \mathbb{O}, *, \circ, b)$ is an intuitionistic fuzzy b -metric space.

Definition 2.5. [19] Let Ω and Ξ be non-void sets and $\mu : \Omega \times \Xi \rightarrow [0, +\infty)$ be a function, such that

- (1) $\mu(\aleph, \varpi) = 0$ if and only if $\aleph = \varpi$, for all $(\aleph, \varpi) \in \Omega \times \Xi$;
- (2) $\mu(\aleph, \varpi) = \mu(\varpi, \aleph)$, for all $(\aleph, \varpi) \in \Omega \cap \Xi$;

(3) $\mu(\aleph, \varpi) \leq \mu(\aleph, \varpi_0) + \mu(\aleph_1, \varpi_0) + \mu(\aleph_1, \varpi)$, for all $\aleph, \aleph_1 \in \Omega$ and $\varpi_0, \varpi \in \Xi$.

The pair (Ω, Ξ, μ) is called a bipolar metric space.

Definition 2.6. [19] Let Ω and Ξ be two non-void sets. We say that the quadruple $(\Omega, \Xi, \mathbb{H}, *)$ is a fuzzy bipolar metric space if $*$ is a continuous t -norm and $\mathbb{H}$ is a fuzzy set on $\Omega \times \Xi \times (0, \infty)$, fulfilling the following conditions for all $\rho, \omega, r > 0$:

- (1) $\mathbb{H}(\aleph, \varpi, \varphi) > 0$ for all $(\aleph, \varpi) \in \Omega \times \Xi$;
- (2) $\mathbb{H}(\aleph, \varpi, \varphi) = 1$ if and only if $\aleph = \varpi$ for $\aleph \in \Omega$ and $\varpi \in \Xi$;
- (3) $\mathbb{H}(\aleph, \varpi, \varphi) = \mathbb{H}(\varpi, \aleph, \varphi)$ for all $\aleph, \varpi \in \Omega \cap \Xi$;
- (4) $\mathbb{H}(\aleph_1, \varphi_2, \rho + \omega + r) \geq \mathbb{H}(\aleph_1, \varphi_1, \rho) * \mathbb{H}(\aleph_2, \varphi_1, \omega) * \mathbb{H}(\aleph_2, \varphi_2, r)$ for all $\aleph_1, \aleph_2 \in \Omega$ and $\varphi_1, \varphi_2 \in \Xi$;
- (5) $\mathbb{H}(\aleph, \varpi, \cdot) : [0, \infty) \rightarrow [0, 1]$ is left continuous;
- (6) $\mathbb{H}(\aleph, \varpi, \cdot)$ is non-decreasing for all $\aleph \in \Omega$ and $\varpi \in \Xi$.

Definition 2.7. [14] Let $\Omega \neq \phi$, $\Xi \neq \phi$ be two sets and $*$ be a CTN, $\circ$ be a CTCN, and $\mathbb{H}, \mathbb{O}$ be neutrosophic sets on $\Omega \times \Xi \times (0, +\infty)$, said to be an intuitionistic fuzzy bipolar metric on $\Omega \times \Xi$, if for all $\aleph, \hat{a} \in \Omega$, $\varpi, \hat{s} \in \Xi$ and $\varphi, \zeta, \hat{w} > 0$, the following conditions are satisfied:

- (1) $\mathbb{H}(\aleph, \varpi, \varphi) + \mathbb{O}(\aleph, \varpi, \varphi) \leq 1$;
- (2) $\mathbb{H}(\aleph, \varpi, \varphi) > 0$;
- (3) $\mathbb{H}(\aleph, \varpi, \varphi) = 1$ for all $\varphi > 0$, if and only if $\aleph = \varpi$;
- (4) $\mathbb{H}(\aleph, \varpi, \varphi) = \mathbb{H}(\varpi, \aleph, \varphi)$;
- (5) $\mathbb{H}(\aleph, \hat{s}, \varphi + \hat{w} + \zeta) \geq \mathbb{H}(\aleph, \varpi, \varphi) * \mathbb{H}(\hat{a}, \varpi, \hat{w}) * \mathbb{H}(\hat{a}, \hat{s}, \zeta)$;
- (6) $\mathbb{H}(\aleph, \varpi, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is continuous and $\lim_{\varphi \rightarrow +\infty} \mathbb{H}(\aleph, \varpi, \varphi) = 1$;
- (7) $\mathbb{H}(\aleph, \varpi, \cdot)$ is an increasing function;
- (8) $\mathbb{O}(\aleph, \varpi, \varphi) < 1$;
- (9) $\mathbb{O}(\aleph, \varpi, \varphi) = 0$ for all $\varphi > 0$, if and only if $\aleph = \varpi$;
- (10) $\mathbb{O}(\aleph, \varpi, \varphi) = \mathbb{O}(\varpi, \aleph, \varphi)$;
- (11) $\mathbb{O}(\aleph, \hat{s}, \varphi + \hat{w} + \zeta) \leq \mathbb{O}(\aleph, \varpi, \varphi) \circ \mathbb{O}(\hat{a}, \varpi, \hat{w}) \circ \mathbb{O}(\hat{a}, \hat{s}, \zeta)$;
- (12) $\mathbb{O}(\aleph, \varpi, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is continuous and $\lim_{\varphi \rightarrow +\infty} \mathbb{O}(\aleph, \varpi, \varphi) = 0$;
- (13) $\mathbb{O}(\aleph, \varpi, \cdot)$ is a decreasing function;
- (14) If $\varphi \leq 0$, then $\mathbb{H}(\aleph, \varpi, \varphi) = 0$ and $\mathbb{O}(\aleph, \varpi, \varphi) = 1$.

Then, $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is called an intuitionistic fuzzy bipolar metric space.

3. FIXED POINT THEOREMS ON NEUTROSOPHIC FUZZY BIPOLAR METRIC SPACE

Definition 3.1. Let $\Omega \neq \phi$, $\Xi \neq \phi$ be two sets and $*$ be a CTN, $\circ$ be a CTCN, and $\mathbb{M} = \{< a, \mathbb{H}(a), \mathbb{O}(a), \mathbb{S}(a) > : a \in \Omega \cap \Xi\}$ be neutrosophic set on $\Omega \times \Xi \times (0, +\infty)$, said to be a neutrosophic fuzzy bipolar metric on $\Omega \times \Xi$, if for all $\aleph, \hat{a} \in \Omega$, $\varpi, \hat{s} \in \Xi$ and $\varphi, \zeta, \hat{w} > 0$, the following conditions are satisfied:

- (1) $\mathbb{H}(\aleph, \varpi, \varphi) + \mathbb{O}(\aleph, \varpi, \varphi) + \mathbb{S}(\aleph, \varpi, \varphi) \leq 3$;

- (2) $\mathbb{H}(\aleph, \varpi, \varphi) > 0$;
- (3) $\mathbb{H}(\aleph, \varpi, \varphi) = 1$ for all $\varphi > 0$, if and only if $\aleph = \varpi$;
- (4) $\mathbb{H}(\aleph, \varpi, \varphi) = \mathbb{H}(\varpi, \aleph, \varphi)$;
- (5) $\mathbb{H}(\aleph, \hat{s}, \varphi + \hat{w} + \zeta) \geq \mathbb{H}(\aleph, \varpi, \varphi) * \mathbb{H}(\hat{a}, \varpi, \hat{w}) * \mathbb{H}(\hat{a}, \hat{s}, \zeta)$;
- (6) $\mathbb{H}(\aleph, \varpi, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is continuous and $\lim_{\varphi \rightarrow +\infty} \mathbb{H}(\aleph, \varpi, \varphi) = 1$;
- (7) $\mathbb{H}(\aleph, \varpi, \cdot)$ is an increasing function;
- (8) $\mathbb{O}(\aleph, \varpi, \varphi) < 1$;
- (9) $\mathbb{O}(\aleph, \varpi, \varphi) = 0$ for all $\varphi > 0$, if and only if $\aleph = \varpi$;
- (10) $\mathbb{O}(\aleph, \varpi, \varphi) = \mathbb{O}(\varpi, \aleph, \varphi)$;
- (11) $\mathbb{O}(\aleph, \hat{s}, \varphi + \hat{w} + \zeta) \leq \mathbb{O}(\aleph, \varpi, \varphi) \circ \mathbb{O}(\hat{a}, \varpi, \hat{w}) \circ \mathbb{O}(\hat{a}, \hat{s}, \zeta)$;
- (12) $\mathbb{O}(\aleph, \varpi, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is continuous and $\lim_{\varphi \rightarrow +\infty} \mathbb{O}(\aleph, \varpi, \varphi) = 0$;
- (13) $\mathbb{O}(\aleph, \varpi, \cdot)$ is a decreasing function;
- (14) $\mathbb{S}(\aleph, \varpi, \varphi) < 1$;
- (15) $\mathbb{S}(\aleph, \varpi, \varphi) = 0$ for all $\varphi > 0$, if and only if $\aleph = \varpi$;
- (16) $\mathbb{S}(\aleph, \varpi, \varphi) = \mathbb{S}(\varpi, \aleph, \varphi)$;
- (17) $\mathbb{S}(\aleph, \hat{s}, \varphi + \hat{w} + \zeta) \leq \mathbb{S}(\aleph, \varpi, \varphi) \circ \mathbb{S}(\hat{a}, \varpi, \hat{w}) \circ \mathbb{S}(\hat{a}, \hat{s}, \zeta)$;
- (18) $\mathbb{S}(\aleph, \varpi, \cdot) : (0, +\infty) \rightarrow [0, 1]$ is continuous and $\lim_{\varphi \rightarrow +\infty} \mathbb{S}(\aleph, \varpi, \varphi) = 0$;
- (19) $\mathbb{S}(\aleph, \varpi, \cdot)$ is a decreasing function;
- (20) If $\varphi \leq 0$, then $\mathbb{H}(\aleph, \varpi, \varphi) = 0$, $\mathbb{O}(\aleph, \varpi, \varphi) = 1$, and $\mathbb{S}(\aleph, \varpi, \varphi) = 1$.

Then, $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is called an neutrosophic fuzzy bipolar metric space.

Example 3.2. Let $\Omega = \{5, 7, 9, 11\}$ and $\Xi = \{4, 6, 8, 10\}$. Define $\mathbb{M} = \{< a, \mathbb{H}(a), \mathbb{O}(a), \mathbb{S}(a) > : a \in \Omega \cap \Xi\}$ in short $\mathbb{M} = (\mathbb{H}, \mathbb{O}, \mathbb{S})$ are neutrosophic set from $\Omega \times \Xi \times (0, +\infty) \rightarrow [0, 1]$ as

$$\begin{aligned}\mathbb{H}(\aleph, \varpi, \varphi) &= \begin{cases} 1, & \text{if } \aleph = \varpi \\ \frac{\varphi}{\varphi + \max\{\aleph, \varpi\}}, & \text{otherwise;} \end{cases} \\ \mathbb{O}(\aleph, \varpi, \varphi) &= \begin{cases} 0, & \text{if } \aleph = \varpi \\ \frac{\max\{\aleph, \varpi\}}{\max\{\aleph, \varpi\} + \varphi}, & \text{otherwise;} \end{cases} \\ \mathbb{S}(\aleph, \varpi, \varphi) &= \begin{cases} 0, & \text{if } \aleph = \varpi \\ \frac{\max\{\aleph, \varpi\}}{\varphi}, & \text{otherwise.} \end{cases}\end{aligned}$$

Then, $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is an neutrosophic fuzzy bipolar metric space with CTN, $\nu * \mu = \nu \mu$ and CTCN, $\nu \circ \mu = \max\{\nu, \mu\}$.

Proof. Here, we prove properties 3.1(5), 3.1(11) and 3.1(17) of Definition 3.1, others are obvious. Let $\aleph = 9$, $\varpi = 6$, $\hat{a} = 7$ and $\hat{s} = 8$. Then,

$$\mathbb{H}(9, 8, \varphi + \hat{w} + \zeta) = \frac{\varphi + \hat{w} + \zeta}{\varphi + \hat{w} + \zeta + \max\{9, 8\}} = \frac{\varphi + \hat{w} + \zeta}{\varphi + \hat{w} + \zeta + 9}.$$

On the other hand,

$$\mathbb{H}(9, 6, \varphi) = \frac{\varphi}{\varphi + \max\{9, 6\}} = \frac{\varphi}{\varphi + 9},$$

$$\mathbb{H}(6, 7, \hat{w}) = \frac{\hat{w}}{\hat{w} + \max\{6, 7\}} = \frac{\hat{w}}{\hat{w} + 7}$$

and

$$\mathbb{H}(7, 8, \zeta) = \frac{\zeta}{\zeta + \max\{7, 8\}} = \frac{\zeta}{\zeta + 8}.$$

Therefore,

$$\frac{\varphi + \hat{w} + \zeta}{\varphi + \hat{w} + \zeta + 9} \geq \frac{\varphi}{\varphi + 9} \cdot \frac{\hat{w}}{\hat{w} + 7} \cdot \frac{\zeta}{\zeta + 8}.$$

Hence,

$$\mathbb{H}(\aleph, \hat{s}, \varphi + \hat{w} + \zeta) \geq \mathbb{H}(\aleph, \varpi, \varphi) * \mathbb{H}(\hat{a}, \varpi, \hat{w}) * \mathbb{H}(\hat{a}, \hat{s}, \zeta), \text{ for all } \varphi, \hat{w}, \zeta > 0.$$

Now,

$$\mathbb{O}(9, 8, \varphi + \hat{w} + \zeta) = \frac{\max\{9, 8\}}{\varphi + \hat{w} + \zeta + \max\{9, 8\}} = \frac{9}{\varphi + \hat{w} + \zeta + 9}.$$

We take,

$$\mathbb{O}(9, 6, \varphi) = \frac{\max\{9, 6\}}{\varphi + \max\{9, 6\}} = \frac{9}{\varphi + 9},$$

$$\mathbb{O}(6, 7, \hat{w}) = \frac{\max\{6, 7\}}{\hat{w} + \max\{6, 7\}} = \frac{7}{\hat{w} + 7}$$

and

$$\mathbb{O}(7, 8, \zeta) = \frac{\max\{7, 8\}}{\zeta + \max\{7, 8\}} = \frac{8}{\zeta + 8}.$$

That is,

$$\frac{9}{\varphi + \hat{w} + \zeta + 9} \leq \max \left\{ \frac{9}{\varphi + 9}, \frac{7}{\hat{w} + 7}, \frac{8}{\zeta + 8} \right\}.$$

Therefore, $\mathbb{O}(\aleph, \hat{s}, \varphi + \hat{w} + \zeta) \leq \mathbb{O}(\aleph, \varpi, \varphi) \circ \mathbb{O}(\hat{a}, \varpi, \hat{w}) \circ \mathbb{O}(\hat{a}, \hat{s}, \zeta)$, for all $\varphi, \hat{w}, \zeta > 0$.

Now,

$$\mathbb{S}(9, 8, \varphi + \hat{w} + \zeta) = \frac{\max\{9, 8\}}{\varphi + \hat{w} + \zeta} = \frac{9}{\varphi + \hat{w} + \zeta}.$$

On the other hand,

$$\mathbb{S}(9, 6, \varphi) = \frac{\max\{9, 6\}}{\varphi} = \frac{9}{\varphi},$$

$$\mathbb{S}(6, 7, \hat{w}) = \frac{\max\{6, 7\}}{\hat{w}} = \frac{7}{\hat{w}}$$

and

$$\mathbb{S}(7, 8, \zeta) = \frac{\max\{7, 8\}}{\zeta} = \frac{8}{\zeta}.$$

That is,

$$\frac{9}{\varphi + \hat{w} + \zeta} \leq \max \left\{ \frac{9}{\varphi}, \frac{7}{\hat{w}}, \frac{8}{\zeta} \right\}.$$

Therefore, $\mathbb{S}(\aleph, \hat{s}, \varphi + \hat{w} + \zeta) \leq \mathbb{S}(\aleph, \varpi, \varphi) \circ \mathbb{S}(\hat{a}, \varpi, \hat{w}) \circ \mathbb{S}(\hat{a}, \hat{s}, \zeta)$, for all $\varphi, \hat{w}, \zeta > 0$.

Hence, $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is an neutrosophic fuzzy bipolar metric space. $\square$

Definition 3.3. Let $\mathcal{P} : \Omega_1 \cup \Xi_1 \rightarrow \Omega_2 \cup \Xi_2$ be a mapping, where (Ω_1, Ξ_1) and (Ω_2, Ξ_2) are pairs of sets. If $\mathcal{P}(\Omega_1) \subseteq \Omega_2$ and $\mathcal{P}(\Xi_1) \subseteq \Xi_2$, then $\mathcal{P}$ is called a covariant map, and this is written as $\mathcal{P} : (\Omega_1, \Xi_1, \mathbb{M}_1, *, \circ) \rightrightarrows (\Omega_2, \Xi_2, \mathbb{M}_2, *, \circ)$, where $\mathbb{M}_1 = (\mathbb{H}_1, \mathbb{O}_1, \mathbb{S}_1)$ and $\mathbb{M}_2 = (\mathbb{H}_2, \mathbb{O}_2, \mathbb{S}_2)$.

Example 3.4. If $\Omega_1 \cup \Xi_1 = X = \{0, 1\}$, then $\mathcal{P}(X) = \{\{\}, \{0\}, \{1\}, X\}$. Suppose $\mathbb{H}(0) = \{\}$ and $\mathbb{H}(1) = X$. Then $\mathcal{P}(\mathbb{H})$ is the function which sends any subset U of X to its image $\mathbb{H}(U)$, which in this case means $\{\} \rightarrow \mathbb{H}(\{\}) = \{\}$, where $\rightarrow$ denotes the mapping under $\mathcal{P}(\mathbb{H})$, so this could also be written as $(\mathcal{P}(\mathbb{H}))(\{\}) = \{\}$. For the other values, $\{0\} \rightarrow \mathbb{H}(\{0\}) = \{\mathbb{H}(0)\} = \{\{\}\}$, $\{1\} \rightarrow \mathbb{H}(\{1\}) = \{\mathbb{H}(1)\} = \{X\}$, $\{0, 1\} \rightarrow \mathbb{H}(\{0, 1\}) = \{\mathbb{H}(0), \mathbb{H}(1)\} = \{\{\}, X\}$.

Definition 3.5. Let $(\Omega, \Xi, \mathbb{M}, *, \circ)$ be an neutrosophic fuzzy bipolar metric space.

- (1) Within the set $\Omega \cup \Xi$, a point $\aleph$ is categorized as a left point if it belongs to Ω , and right point if it belongs to Ξ , and a central point if it belongs to both.
- (2) A sequence denoted as $\{\aleph_\alpha\}$ and belonging to the set Ω is termed a left sequence, while a sequence denoted as $\{\alpha_n\}$ and belonging to the set Ξ is referred to as a right sequence.
- (3) A sequence denoted as $\{\aleph_\alpha\} \subset \Omega \cup \Xi$ is considered to converge to a point $\aleph$ if and only if $\{\aleph_\alpha\}$ is a left sequence, $\aleph$ is a right point and

$$\begin{aligned} \lim_{\alpha \rightarrow +\infty} \mathbb{H}(\aleph_\alpha, \aleph, \varphi) &= 1, \\ \lim_{\alpha \rightarrow +\infty} \mathbb{O}(\aleph_\alpha, \aleph, \varphi) &= 0, \\ \lim_{\alpha \rightarrow +\infty} \mathbb{S}(\aleph_\alpha, \aleph, \varphi) &= 0, \end{aligned}$$

for all $\varphi > 0$.

- (4) A sequence $\{(\aleph_\alpha, \alpha_\alpha)\} \subset \Omega \times \Xi$ is called a bisequence. If both sequences $\{\aleph_\alpha\}$ and $\{\alpha_\alpha\}$ converge, then the bisequence $\{(\aleph_\alpha, \alpha_\alpha)\}$ is called convergent within the space $\Omega \times \Xi$.
- (5) If $\{\aleph_\alpha\}$ and $\{\alpha_\alpha\}$ both converge to a common point $\alpha \in \Omega \cap \Xi$, then the bisequence $\{(\aleph_\alpha, \alpha_\alpha)\}$ is labeled as biconvergent. A sequence $\{(\aleph_\alpha, \alpha_\alpha)\}$ is a Cauchy bisequence if

$$\begin{aligned} \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{H}(\aleph_\alpha, \alpha_\kappa, \varphi) &= 1, \\ \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{O}(\aleph_\alpha, \alpha_\kappa, \varphi) &= 0, \\ \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{S}(\aleph_\alpha, \alpha_\kappa, \varphi) &= 0, \end{aligned}$$

for all $\varphi > 0$.

- (6) An neutrosophic fuzzy bipolar metric space is said to be complete if every Cauchy bisequence is convergent.

Example 3.6. Let $\Omega = (1, \infty)$ and $\Xi = [-1, 1]$. Define $\mathbb{H} : \Omega \times \Xi \rightarrow \mathbf{R}^+$ as $\mathbb{H}(\aleph, \varpi) = |\aleph^2 - \varpi^2|$. Then $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is an neutrosophic fuzzy bipolar metric space.

Note that the left sequence $(1 + \frac{1}{\alpha})$ converges to right points 1 and -1 .

Lemma 3.7. Let $\{\aleph_\alpha\}$ be a Cauchy sequence in the neutrosophic fuzzy bipolar metric space $(\Omega, \Xi, \mathbb{M}, *, \circ)$ such that $\aleph_\alpha \neq \aleph_\kappa$, whenever $\kappa, \alpha \in \mathbf{N}$ with $\alpha \neq \kappa$. Then the sequence $\{\aleph_\alpha\}$ can converge to, at most, one limit point.

Proof. Assume that, $\aleph_\alpha \rightarrow \aleph \in \Xi$ and $\aleph_\alpha \rightarrow \varpi \in \Omega \cap \Xi$, for $\aleph \neq \varpi$. Then,

$$\lim_{\alpha \rightarrow +\infty} \mathbb{H}(\aleph_\alpha, \aleph, \varphi) = 1, \lim_{\alpha \rightarrow +\infty} \mathbb{O}(\aleph_\alpha, \aleph, \varphi) = 0, \lim_{\alpha \rightarrow +\infty} \mathbb{S}(\aleph_\alpha, \aleph, \varphi) = 0$$

and

$$\lim_{\alpha \rightarrow +\infty} \mathbb{H}(\aleph_\alpha, \varpi, \varphi) = 1, \lim_{\alpha \rightarrow +\infty} \mathbb{O}(\aleph_\alpha, \varpi, \varphi) = 0, \lim_{\alpha \rightarrow +\infty} \mathbb{S}(\aleph_\alpha, \varpi, \varphi) = 0,$$

for all $\varphi > 0$.

Suppose,

$$\begin{aligned} \mathbb{H}(\aleph, \varpi, \varphi) &\geq \mathbb{H}\left(\aleph, \aleph_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_\alpha, \aleph_{\alpha+1}, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi, \frac{\varphi}{3}\right) \\ &\rightarrow 1 * 1 * 1, \text{ as } \alpha \rightarrow +\infty, \\ \mathbb{O}(\aleph, \varpi, \varphi) &\leq \mathbb{O}\left(\aleph, \aleph_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_\alpha, \aleph_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi, \frac{\varphi}{3}\right) \\ &\rightarrow 0 \circ 0 \circ 0, \text{ as } \alpha \rightarrow +\infty, \\ \mathbb{S}(\aleph, \varpi, \varphi) &\leq \mathbb{S}\left(\aleph, \aleph_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_\alpha, \aleph_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi, \frac{\varphi}{3}\right) \\ &\rightarrow 0 \circ 0 \circ 0, \text{ as } \alpha \rightarrow +\infty. \end{aligned}$$

That is,

$$\mathbb{H}(\aleph, \varpi, \varphi) \geq 1 * 1 * 1 = 1, \mathbb{O}(\aleph, \varpi, \varphi) \leq 0 \circ 0 \circ 0 = 0, \mathbb{S}(\aleph, \varpi, \varphi) \leq 0 \circ 0 \circ 0 = 0.$$

Hence $\aleph = \varpi$. □

Lemma 3.8. Let $(\Omega, \Xi, \mathbb{M}, *, \circ)$ be an neutrosophic fuzzy bipolar metric space. If for some $0 < \theta < 1$ and for any $\aleph, \varpi \in \Omega$, $\varphi > 0$,

$$\mathbb{H}(\aleph, \varpi, \varphi) \geq \mathbb{H}\left(\aleph, \varpi, \frac{\varphi}{\theta}\right), \mathbb{O}(\aleph, \varpi, \varphi) \leq \mathbb{O}\left(\aleph, \varpi, \frac{\varphi}{\theta}\right), \mathbb{S}(\aleph, \varpi, \varphi) \leq \mathbb{S}\left(\aleph, \varpi, \frac{\varphi}{\theta}\right), \quad (1)$$

then $\aleph = \varpi$.

Proof. Equation (1) implies that,

$$\mathbb{H}(\aleph, \varpi, \varphi) \geq \mathbb{H}(\aleph, \varpi, \frac{\varphi}{\theta^\alpha}), \quad \mathbb{O}(\aleph, \varpi, \varphi) \leq \mathbb{O}(\aleph, \varpi, \frac{\varphi}{\theta^\alpha}) \text{ and}$$

$$\mathbb{S}(\aleph, \varpi, \varphi) \leq \mathbb{S}(\aleph, \varpi, \frac{\varphi}{\theta^\alpha}), \alpha \in \mathbf{N}, \varphi > 0.$$

Now,

$$\mathbb{H}(\aleph, \varpi, \varphi) \geq \lim_{\alpha \rightarrow +\infty} \mathbb{H}(\aleph, \varpi, \frac{\varphi}{\theta^\alpha}) = 1,$$

$$\mathbb{O}(\aleph, \varpi, \varphi) \leq \lim_{\alpha \rightarrow +\infty} \mathbb{O}(\aleph, \varpi, \frac{\varphi}{\theta^\alpha}) = 0,$$

$$\mathbb{S}(\aleph, \varpi, \varphi) \leq \lim_{\alpha \rightarrow +\infty} \mathbb{S}(\aleph, \varpi, \frac{\varphi}{\theta^\alpha}) = 0.$$

Also, by the Definition 3.1, we get $\aleph = \varpi$. □

Theorem 3.9. Suppose $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is a complete neutrosophic fuzzy bipolar metric space with $0 < \theta < 1$. Let $\mathcal{P} : \Omega \cup \Xi \rightarrow \Omega \cup \Xi$ be a mapping satisfying:

- (1) $\mathcal{P}(\Omega) \subseteq \Omega$ and $\mathcal{P}(\Xi) \subseteq \Xi$;
- (2) $\mathbb{H}(\mathcal{P}\aleph, \mathcal{P}\varpi, \theta\varphi) \geq \mathbb{H}(\aleph, \varpi, \varphi)$, $\mathbb{O}(\mathcal{P}\aleph, \mathcal{P}\varpi, \theta\varphi) \leq \mathbb{O}(\aleph, \varpi, \varphi)$, $\mathbb{S}(\mathcal{P}\aleph, \mathcal{P}\varpi, \theta\varphi) \leq \mathbb{S}(\aleph, \varpi, \varphi)$, for all $\aleph \in \Omega$, $\varpi \in \Xi$, and $\varphi > 0$.

Then $\mathcal{P}$ has a unique fixed point.

Proof. Let $\aleph_0 \in \Omega$ and $\varpi_0 \in \Xi$ and assume that $\mathcal{P}(\aleph_\alpha) = \aleph_{\alpha+1}$ and $\mathcal{P}(\varpi_\alpha) = \varpi_{\alpha+1}$ for all $\alpha \in \mathbf{N} \cup \{0\}$.

We get, $(\aleph_\alpha, \varpi_\alpha)$ as a bisequence on the neutrosophic fuzzy bipolar metric space $(\Omega, \Xi, \mathbb{M}, *, \circ)$. Now, we have

$$\mathbb{H}(\aleph_1, \varpi_1, \varphi) = \mathbb{H}(\mathcal{P}\aleph_0, \mathcal{P}\varpi_0, \varphi) \geq \mathbb{H}(\aleph_0, \varpi_0, \frac{\varphi}{\theta}),$$

$$\mathbb{O}(\aleph_1, \varpi_1, \varphi) = \mathbb{O}(\mathcal{P}\aleph_0, \mathcal{P}\varpi_0, \varphi) \leq \mathbb{O}(\aleph_0, \varpi_0, \frac{\varphi}{\theta})$$

and

$$\mathbb{S}(\aleph_1, \varpi_1, \varphi) = \mathbb{S}(\mathcal{P}\aleph_0, \mathcal{P}\varpi_0, \varphi) \leq \mathbb{S}(\aleph_0, \varpi_0, \frac{\varphi}{\theta}),$$

for all $\varphi > 0$ and $\alpha \in \mathbf{N}$. By simple induction, we get

$$\begin{aligned} \mathbb{H}(\aleph_\alpha, \varpi_\alpha, \varphi) &= \mathbb{H}(\mathcal{P}\aleph_{\alpha-1}, \mathcal{P}\varpi_{\alpha-1}, \varphi) \geq \mathbb{H}(\aleph_{\alpha-1}, \varpi_{\alpha-1}, \frac{\varphi}{\theta}) \\ &\geq \mathbb{H}(\aleph_{\alpha-2}, \varpi_{\alpha-2}, \frac{\varphi}{\theta^2}) \geq \mathbb{H}(\aleph_{\alpha-3}, \varpi_{\alpha-3}, \frac{\varphi}{\theta^3}) \geq \dots \geq \mathbb{H}(\aleph_0, \varpi_0, \frac{\varphi}{\theta^\alpha}), \end{aligned}$$

$$\begin{aligned} \mathbb{O}(\aleph_\alpha, \varpi_\alpha, \varphi) &= \mathbb{O}(\mathcal{P}\aleph_{\alpha-1}, \mathcal{P}\varpi_{\alpha-1}, \varphi) \leq \mathbb{O}(\aleph_{\alpha-1}, \varpi_{\alpha-1}, \frac{\varphi}{\theta}) \\ &\leq \mathbb{O}(\aleph_{\alpha-2}, \varpi_{\alpha-2}, \frac{\varphi}{\theta^2}) \leq \mathbb{O}(\aleph_{\alpha-3}, \varpi_{\alpha-3}, \frac{\varphi}{\theta^3}) \leq \dots \leq \mathbb{O}(\aleph_0, \varpi_0, \frac{\varphi}{\theta^\alpha}) \end{aligned}$$

and

$$\mathbb{S}(\aleph_\alpha, \varpi_\alpha, \varphi) = \mathbb{S}(\mathcal{P}\aleph_{\alpha-1}, \mathcal{P}\varpi_{\alpha-1}, \varphi) \leq \mathbb{S}(\aleph_{\alpha-1}, \varpi_{\alpha-1}, \frac{\varphi}{\theta})$$

$$\leq \mathbb{S}\left(\aleph_{\alpha-2}, \varpi_{\alpha-2}, \frac{\varphi}{\theta^2}\right) \leq \mathbb{S}\left(\aleph_{\alpha-3}, \varpi_{\alpha-3}, \frac{\varphi}{\theta^3}\right) \leq \dots \leq \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{\theta^\alpha}\right).$$

We obtain,

$$\mathbb{H}(\aleph_\alpha, \varpi_\alpha, \varphi) \geq \mathbb{H}(\aleph_0, \varpi_0, \frac{\varphi}{\theta^\alpha}), \quad \mathbb{O}(\aleph_\alpha, \varpi_\alpha, \varphi) \leq \mathbb{O}(\aleph_0, \varpi_0, \frac{\varphi}{\theta^\alpha}),$$

$$\mathbb{S}(\aleph_\alpha, \varpi_\alpha, \varphi) \leq \mathbb{S}(\aleph_0, \varpi_0, \frac{\varphi}{\theta^\alpha})$$

and

$$\mathbb{H}(\aleph_{\alpha+1}, \varpi_\alpha, \varphi) \geq \mathbb{H}(\aleph_1, \varpi_0, \frac{\varphi}{\theta^\alpha}), \quad \mathbb{O}(\aleph_{\alpha+1}, \varpi_\alpha, \varphi) \leq \mathbb{O}(\aleph_1, \varpi_0, \frac{\varphi}{\theta^\alpha}),$$

$$\mathbb{S}(\aleph_{\alpha+1}, \varpi_\alpha, \varphi) \leq \mathbb{S}(\aleph_1, \varpi_0, \frac{\varphi}{\theta^\alpha}).$$

Letting, $\alpha < \kappa$, for $\alpha, \kappa \in \mathbb{N}$. Then,

$$\begin{aligned} \mathbb{H}(\aleph_\alpha, \varpi_\kappa, \varphi) &\geq \mathbb{H}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\kappa, \frac{\varphi}{3}\right) * \dots \\ &\geq \mathbb{H}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \dots * \\ &\quad \mathbb{H}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{H}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{H}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right), \\ \mathbb{O}(\aleph_\alpha, \varpi_\kappa, \varphi) &\leq \mathbb{O}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\kappa, \frac{\varphi}{3}\right) \circ \dots \\ &\leq \mathbb{O}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \dots \circ \\ &\quad \mathbb{O}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \mathbb{O}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \mathbb{O}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right) \end{aligned}$$

and

$$\begin{aligned} \mathbb{S}(\aleph_\alpha, \varpi_\kappa, \varphi) &\leq \mathbb{S}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\kappa, \frac{\varphi}{3}\right) \circ \dots \\ &\leq \mathbb{S}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \dots \circ \\ &\quad \mathbb{S}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \mathbb{S}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \mathbb{S}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right). \end{aligned}$$

Therefore,

$$\begin{aligned} \mathbb{H}(\aleph_\alpha, \varpi_\kappa, \varphi) &\geq \mathbb{H}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \dots * \mathbb{H}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \\ &\quad \mathbb{H}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{H}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right) \geq \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) * \\ &\quad \mathbb{H}\left(\aleph_1, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) * \dots * \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) * \mathbb{H}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) * \\ &\quad \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^\kappa}\right), \\ \mathbb{O}(\aleph_\alpha, \varpi_\kappa, \varphi) &\leq \mathbb{O}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \dots \circ \mathbb{O}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \\ &\quad \mathbb{O}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{O}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right) \leq \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) \circ \end{aligned}$$

$$\begin{aligned} & \mathbb{O}\left(\aleph_1, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) \circ \cdots \circ \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) \circ \mathbb{O}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) \circ \\ & \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^\kappa}\right) \end{aligned}$$

and

$$\begin{aligned} \mathbb{S}(\aleph_\alpha, \varpi_\kappa, \varphi) & \leq \mathbb{S}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \cdots \circ \mathbb{S}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \\ & \mathbb{S}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{S}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right) \\ & \leq \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) \circ \mathbb{S}\left(\aleph_1, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) \circ \cdots \circ \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) \circ \\ & \mathbb{S}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) \circ \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^\kappa}\right), \end{aligned}$$

implies that,

$$\begin{aligned} \mathbb{H}(\aleph_\alpha, \varpi_\kappa, \varphi) & \geq \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) * \mathbb{H}\left(\aleph_1, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) * \cdots * \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) * \\ & \mathbb{H}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) * \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^\kappa}\right), \\ \mathbb{O}(\aleph_\alpha, \varpi_\kappa, \varphi) & \leq \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) \circ \mathbb{O}\left(\aleph_1, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) \circ \cdots \circ \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) \circ \\ & \mathbb{O}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) \circ \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^\kappa}\right) \end{aligned}$$

and

$$\begin{aligned} \mathbb{S}(\aleph_\alpha, \varpi_\kappa, \varphi) & \leq \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) \circ \mathbb{S}\left(\aleph_1, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) \circ \cdots \circ \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) \circ \\ & \mathbb{S}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^{\kappa-1}}\right) \circ \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}\theta^\kappa}\right). \end{aligned}$$

As $\alpha, \kappa \rightarrow +\infty$, we deduce

$$\begin{aligned} \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{H}(\aleph_\alpha, \varpi_\kappa, \varphi) & = 1 * 1 * \cdots * 1 = 1, \\ \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{O}(\aleph_\alpha, \varpi_\kappa, \varphi) & = 0 \circ 0 \circ \cdots \circ 0 = 0, \end{aligned}$$

$$\text{and } \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{S}(\aleph_\alpha, \varpi_\kappa, \varphi) = 0 \circ 0 \circ \cdots \circ 0 = 0.$$

Which implies that the bisequence $(\aleph_\alpha, \varpi_\alpha)$ is a Cauchy bisequence. Since $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is a complete neutrosophic fuzzy bipolar metric space, we have that $\{\aleph_\alpha\} \rightarrow \aleph$ and $\{\varpi_\alpha\} \rightarrow \varpi$, where $\aleph \in \Omega \cap \Xi$. Using Lemmas 3.7 and 3.8, we get

$$\begin{aligned} \mathbb{H}(\aleph, \mathcal{P}\aleph, \varphi) & \geq \mathbb{H}\left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \aleph_{\alpha+1}, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3}\right) \\ & = \mathbb{H}\left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3}\right) * \mathbb{H}\left(\mathcal{P}\aleph_\alpha, \mathcal{P}\aleph_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\mathcal{P}\aleph_\alpha, \mathcal{P}\aleph, \frac{\varphi}{3}\right) \\ & \rightarrow 1 * 1 * 1 = 1 \quad \text{as } \alpha \rightarrow +\infty, \\ \mathbb{O}(\aleph, \mathcal{P}\aleph, \varphi) & \leq \mathbb{O}\left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \aleph_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3}\right) \end{aligned}$$

$$\begin{aligned}
&= \mathbb{O} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{O} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{O} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\
&\rightarrow 0 \circ 0 \circ 0 = 0 \quad \text{as } \alpha \rightarrow +\infty \quad \text{and} \\
\mathbb{S}(\aleph, \mathcal{P}\aleph, \varphi) &\leq \mathbb{S} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\aleph_{\alpha+1}, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\
&= \mathbb{S} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\
&\rightarrow 0 \circ 0 \circ 0 = 0 \quad \text{as } \alpha \rightarrow +\infty.
\end{aligned}$$

Hence, $\mathcal{P}\aleph = \aleph$.

Now, we examine the uniqueness. Let $\mathcal{P}\omega = \omega$, for some $\omega \in \Omega \cap \Xi$, then

$$\begin{aligned}
1 &\geq \mathbb{H}(\omega, \aleph, \varphi) = \mathbb{H}(\mathcal{P}\omega, \mathcal{P}\aleph, \varphi) \geq \mathbb{H} \left(\omega, \aleph, \frac{\varphi}{\theta} \right) = \mathbb{H} \left(\mathcal{P}\omega, \mathcal{P}\aleph, \frac{\varphi}{\theta} \right) \\
&\geq \mathbb{H} \left(\omega, \aleph, \frac{\varphi}{\theta^2} \right) \geq \cdots \geq \mathbb{H} \left(\omega, \aleph, \frac{\varphi}{\theta^\alpha} \right) \rightarrow 1 \quad \text{as } \alpha \rightarrow +\infty \text{ and} \\
0 &\leq \mathbb{O}(\omega, \aleph, \varphi) = \mathbb{O}(\mathcal{P}\omega, \mathcal{P}\aleph, \varphi) \leq \mathbb{O} \left(\omega, \aleph, \frac{\varphi}{\theta} \right) = \mathbb{O} \left(\mathcal{P}\omega, \mathcal{P}\aleph, \frac{\varphi}{\theta} \right) \\
&\leq \mathbb{O} \left(\omega, \aleph, \frac{\varphi}{\theta^2} \right) \leq \cdots \leq \mathbb{O} \left(\omega, \aleph, \frac{\varphi}{\theta^\alpha} \right) \rightarrow 0 \quad \text{as } \alpha \rightarrow +\infty, \\
0 &\leq \mathbb{S}(\omega, \aleph, \varphi) = \mathbb{S}(\mathcal{P}\omega, \mathcal{P}\aleph, \varphi) \leq \mathbb{S} \left(\omega, \aleph, \frac{\varphi}{\theta} \right) = \mathbb{S} \left(\mathcal{P}\omega, \mathcal{P}\aleph, \frac{\varphi}{\theta} \right) \\
&\leq \mathbb{S} \left(\omega, \aleph, \frac{\varphi}{\theta^2} \right) \leq \cdots \leq \mathbb{S} \left(\omega, \aleph, \frac{\varphi}{\theta^\alpha} \right) \rightarrow 0 \quad \text{as } \alpha \rightarrow +\infty,
\end{aligned}$$

by using Lemmas 3.7 and 3.8, $\aleph = \omega$. □

Theorem 3.10. *Let $(\Omega, \Xi, \mathbb{M}_b, *, \circ)$ be a complete neutrosophic fuzzy bipolar metric space such that*

$$\lim_{\varphi \rightarrow \infty} \mathbb{H}(\aleph, \varpi, \varphi) = 1, \lim_{\varphi \rightarrow \infty} \mathbb{O}(\aleph, \varpi, \varphi) = 0 \text{ and } \lim_{\varphi \rightarrow \infty} \mathbb{S}(\aleph, \varpi, \varphi) = 0,$$

for all $\aleph \in \Omega, \varpi \in \Xi$.

Let $\mathcal{P}, \mathcal{Q} : \Xi \cup \Omega \rightarrow \Xi \cup \Omega$ be two mappings satisfying

- (1) $\mathcal{P}(\Xi) \subseteq \Xi, \mathcal{Q}(\Xi) \subseteq \Xi$, and $\mathcal{P}(\Omega) \subseteq \Omega, \mathcal{Q}(\Omega) \subseteq \Omega$.
- (2) $\mathbb{H}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) \geq \mathbb{H}(\aleph, \varpi, \varphi), \mathbb{O}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) \leq \mathbb{O}(\aleph, \varpi, \varphi)$ and $\mathbb{S}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) \leq \mathbb{S}(\aleph, \varpi, \varphi)$ for all $\aleph \in \Xi, \varpi \in \Omega$ and $\varphi > 0$, where $0 < \theta < 1$

Then, $\mathcal{P}$ and $\mathcal{Q}$ have a unique common fixed point.

Proof. Fix $\aleph_0 \in \Xi$ and $\varpi_0 \in \Omega$ and assume that $\mathcal{P}(\aleph_{2\alpha}) = \aleph_{2\alpha+1}, \mathcal{Q}(\aleph_{2\alpha+1}) = \aleph_{2\alpha+2}, \mathcal{P}(\varpi_{2\alpha}) = \varpi_{2\alpha+1}$ and $\mathcal{Q}(\varpi_{2\alpha+1}) = \varpi_{2\alpha+2}$ for all $\alpha \in \mathbf{N} \cup \{0\}$. Then, we get $(\aleph_\alpha, \varpi_\alpha)$ as a bisequence on the neutrosophic fuzzy bipolar metric space $(\Xi, \Omega, \mathbb{M}, *, \circ)$. Now, we have

$$\begin{aligned}
\mathbb{H}(\aleph_1, \varpi_1, \varphi) &= \mathbb{H}(\mathcal{P}(\aleph_0), \mathcal{Q}(\varpi_0), \varphi) \geq \mathbb{H} \left(\aleph_0, \varpi_0, \frac{\varphi}{\theta} \right), \\
\mathbb{O}(\aleph_1, \varpi_1, \varphi) &= \mathbb{O}(\mathcal{P}(\aleph_0), \mathcal{Q}(\varpi_0), \varphi) \leq \mathbb{O} \left(\aleph_0, \varpi_0, \frac{\varphi}{\theta} \right),
\end{aligned}$$

$$\mathbb{S}(\aleph_1, \varpi_1, \varphi) = \mathbb{S}(\mathcal{P}(\aleph_0), \mathcal{Q}(\varpi_0), \varphi) \leq \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{\theta}\right),$$

for all $\varphi > 0$ and $\alpha \in \mathbf{N}$. By induction, we obtain

$$\begin{aligned}\mathbb{H}(\aleph_{2\alpha+1}, \varpi_{2\alpha+1}, \varphi) &= \mathbb{H}(\mathcal{P}(\aleph_{2\alpha}), \mathcal{Q}(\varpi_{2\alpha}), \varphi) \geq \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{\theta^{2\alpha+1}}\right), \\ \mathbb{H}(\aleph_{2\alpha+1}, \varpi_{2\alpha+2}, \varphi) &= \mathbb{H}(\mathcal{P}(\aleph_{2\alpha}), \mathcal{Q}(\varpi_{2\alpha+1}), \varphi) \geq \mathbb{H}\left(\aleph_0, \varpi_1, \frac{\varphi}{\theta^{2\alpha+1}}\right), \\ \mathbb{O}(\aleph_{2\alpha+1}, \varpi_{2\alpha+1}, \varphi) &= \mathbb{O}(\mathcal{P}(\aleph_{2\alpha}), \mathcal{Q}(\varpi_{2\alpha}), \varphi) \leq \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{\theta^{2\alpha+1}}\right), \\ \mathbb{O}(\aleph_{2\alpha+1}, \varpi_{2\alpha+2}, \varphi) &= \mathbb{O}(\mathcal{P}(\aleph_{2\alpha}), \mathcal{Q}(\varpi_{2\alpha+1}), \varphi) \leq \mathbb{O}\left(\aleph_0, \varpi_1, \frac{\varphi}{\theta^{2\alpha+1}}\right)\end{aligned}$$

and

$$\begin{aligned}\mathbb{S}(\aleph_{2\alpha+1}, \varpi_{2\alpha+1}, \varphi) &= \mathbb{S}(\mathcal{P}(\aleph_{2\alpha}), \mathcal{Q}(\varpi_{2\alpha}), \varphi) \leq \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{\theta^{2\alpha+1}}\right), \\ \mathbb{S}(\aleph_{2\alpha+1}, \varpi_{2\alpha+2}, \varphi) &= \mathbb{S}(\mathcal{P}(\aleph_{2\alpha}), \mathcal{Q}(\varpi_{2\alpha+1}), \varphi) \leq \mathbb{S}\left(\aleph_0, \varpi_1, \frac{\varphi}{\theta^{2\alpha+1}}\right),\end{aligned}$$

for all $\varphi > 0$ and $\alpha \in \mathbf{N}$.

Letting $\alpha < \beta$, for $\alpha, \beta \in \mathbf{N}$. Then, from the definition of the neutrosophic fuzzy bipolar metric space, we get

$$\begin{aligned}\mathbb{H}(\aleph_\alpha, \varpi_\beta, \varphi) &\geq \mathbb{H}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\beta, \frac{\varphi}{3}\right) \dots \\ &\geq \mathbb{H}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \dots \\ &\mathbb{H}\left(\aleph_{\beta-1}, \varpi_{\beta-1}, \frac{\varphi}{3^{\beta-1}}\right) * \mathbb{H}\left(\aleph_\beta, \varpi_{\beta-1}, \frac{\varphi}{3^{\beta-1}}\right) * \mathbb{H}\left(\aleph_{\beta-1}, \varpi_\beta, \frac{\varphi}{3^\beta}\right), \\ \mathbb{O}(\aleph_\alpha, \varpi_\beta, \varphi) &\leq \mathbb{O}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\beta, \frac{\varphi}{3}\right) \dots \\ &\leq \mathbb{O}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \dots \\ &\mathbb{O}\left(\aleph_{\beta-1}, \varpi_{\beta-1}, \frac{\varphi}{3^{\beta-1}}\right) \circ \mathbb{O}\left(\aleph_\beta, \varpi_{\beta-1}, \frac{\varphi}{3^{\beta-1}}\right) \circ \mathbb{O}\left(\aleph_{\beta-1}, \varpi_\beta, \frac{\varphi}{3^\beta}\right)\end{aligned}$$

and

$$\begin{aligned}\mathbb{S}(\aleph_\alpha, \varpi_\beta, \varphi) &\leq \mathbb{S}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\beta, \frac{\varphi}{3}\right) \dots \\ &\leq \mathbb{S}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \dots \\ &\mathbb{S}\left(\aleph_{\beta-1}, \varpi_{\beta-1}, \frac{\varphi}{3^{\beta-1}}\right) \circ \mathbb{S}\left(\aleph_\beta, \varpi_{\beta-1}, \frac{\varphi}{3^{\beta-1}}\right) \circ \mathbb{S}\left(\aleph_{\beta-1}, \varpi_\beta, \frac{\varphi}{3^\beta}\right).\end{aligned}$$

Therefore,

$$\begin{aligned}\mathbb{H}(\aleph_\alpha, \varpi_\beta, \varphi) &\geq \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) * \mathbb{H}\left(\aleph_0, \varpi_1, \frac{\varphi}{3\theta^{\alpha+1}}\right) \dots \mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\beta-1}\theta^{\beta-1}}\right), \\ \mathbb{O}(\aleph_\alpha, \varpi_\beta, \varphi) &\leq \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) * \mathbb{O}\left(\aleph_0, \varpi_1, \frac{\varphi}{3\theta^{\alpha+1}}\right) \dots \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\beta-1}\theta^{\beta-1}}\right)\end{aligned}$$

and

$$\mathbb{S}(\aleph_\alpha, \varpi_\beta, \varphi) \leq \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3\theta^\alpha}\right) * \mathbb{S}\left(\aleph_0, \varpi_1, \frac{\varphi}{3\theta^{\alpha+1}}\right) \dots \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\beta-1}\theta^{\beta-1}}\right).$$

As $\alpha, \beta \rightarrow \infty$, we get

$$\mathbb{H}(\aleph_\alpha, \varpi_\beta, \varphi) \geq 1, \mathbb{O}(\aleph_\alpha, \varpi_\beta, \varphi) \leq 0, \mathbb{S}(\aleph_\alpha, \varpi_\beta, \varphi) \leq 0 \quad \text{for all } \varphi > 0.$$

Thus, the bisequence $(\aleph_\alpha, \varpi_\alpha)$ is a Cauchy bisequence. Since $(\Xi, \Omega, \mathbb{M}, *, \circ)$ is a complete neutrosophic fuzzy bipolar metric space. By Lemma 3.7, the bisequence $(\aleph_\alpha, \varpi_\alpha)$ is a biconvergent sequence. Therefore, $\aleph_\alpha \rightarrow u$ and $\varpi_\alpha \rightarrow u$, where $u \in \Xi \cap \Omega$. By Lemma 3.8, both sequences $\aleph_\alpha$ and ϖ_α have a unique limit. From the triangular property of neutrosophic bipolar metric spaces, we have

$$\begin{aligned} \mathbb{H}(\mathcal{P}u, u, \varphi) &\geq \mathbb{H}\left(\mathcal{P}(u), \varpi_{\alpha+1}, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right) \\ &= \mathbb{H}\left(\mathcal{P}u, \mathcal{Q}(\varpi_\alpha), \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right) \\ &\geq \mathbb{H}\left(u, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right), \\ \mathbb{O}(\mathcal{P}u, u, \varphi) &\leq \mathbb{O}\left(\mathcal{P}(u), \varpi_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right) \\ &= \mathbb{O}\left(\mathcal{P}u, \mathcal{Q}(\varpi_\alpha), \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right) \\ &\leq \mathbb{O}\left(u, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right) \end{aligned}$$

and

$$\begin{aligned} \mathbb{S}(\mathcal{P}u, u, \varphi) &\leq \mathbb{S}\left(\mathcal{P}(u), \varpi_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right) \\ &= \mathbb{S}\left(\mathcal{P}u, \mathcal{Q}(\varpi_\alpha), \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right) \\ &\leq \mathbb{S}\left(u, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_{\alpha+1}, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, u, \frac{\varphi}{3}\right), \end{aligned}$$

for all $\alpha \in \mathbf{N}$ and $\varphi > 0$ and as $\alpha \rightarrow \infty$, we get

$$\mathbb{H}(\mathcal{P}u, u, \varphi) \rightarrow 1 * 1 * 1 = 1, \mathbb{O}(\mathcal{P}u, u, \varphi) \rightarrow 0 \circ 0 \circ 0 = 0, \mathbb{S}(\mathcal{P}u, u, \varphi) \rightarrow 0 \circ 0 \circ 0 = 0.$$

From Definition 3.1 condition (2), $\mathcal{P}(u) = u$. Again,

$$\begin{aligned} \mathbb{H}(u, \mathcal{Q}(u), \varphi) &= \mathbb{H}(\mathcal{P}(u), \mathcal{Q}(u), \varphi) \geq \mathbb{H}\left(u, u, \frac{\varphi}{\theta}\right) = 1, \\ \mathbb{O}(u, \mathcal{Q}(u), \varphi) &= \mathbb{O}(\mathcal{P}(u), \mathcal{Q}(u), \varphi) \leq \mathbb{O}\left(u, u, \frac{\varphi}{\theta}\right) = 0, \\ \mathbb{S}(u, \mathcal{Q}(u), \varphi) &= \mathbb{S}(\mathcal{P}(u), \mathcal{Q}(u), \varphi) \leq \mathbb{S}\left(u, u, \frac{\varphi}{\theta}\right) = 0. \end{aligned}$$

Therefore, $\mathcal{Q}(u) = u$. Hence, u is a common fixed point of $\mathcal{P}$ and $\mathcal{Q}$.

Let $v \in \Xi \cap \Omega$ be another fixed point of $\mathcal{P}$ and $\mathcal{Q}$. Then,

$$\begin{aligned} \mathbb{H}(u, v, \varphi) &= \mathbb{H}(\mathcal{P}(u), \mathcal{Q}(v), \varphi) \geq \mathbb{H}\left(u, v, \frac{\varphi}{\theta}\right), \\ \mathbb{O}(u, v, \varphi) &= \mathbb{O}(\mathcal{P}(u), \mathcal{Q}(v), \varphi) \leq \mathbb{O}\left(u, v, \frac{\varphi}{\theta}\right), \end{aligned}$$

$$\mathbb{S}(u, v, \varphi) = \mathbb{S}\left(\mathcal{P}(u), \mathcal{Q}(v), \varphi\right) \leq \mathbb{S}\left(u, v, \frac{\varphi}{\theta}\right),$$

for $0 < \theta < 1$ and for all $\varphi > 0$. By Lemma 3.8, we have $u = v$. $\square$

Example 3.11. Let $\Xi = [0, 2]$ and $\Omega = \{0\} \cup \mathbf{N} - \{1, 2\}$. Define,

$$\mathbb{H} = \frac{\varphi}{(\varphi + |\aleph - \varpi|)}, \quad \mathbb{O} = \frac{|\aleph - \varpi|}{(\varphi + |\aleph - \varpi|)}, \quad \mathbb{S} = \frac{|\aleph - \varpi|}{\varphi}$$

for all $\varphi > 0, \aleph \in \Xi$ and $\varpi \in \Omega$. Clearly, $(\Xi, \Omega, \mathbb{M}, *, \circ)$ is a complete neutrosophic fuzzy bipolar metric space, where $*$ is a continuous t -norm defined as $p * q = pq$.

Let $\mathcal{P}, \mathcal{Q} : \Xi \cup \Omega \rightarrow \Xi \cup \Omega$ be mappings defined by

$$\mathcal{P}(\aleph) = \begin{cases} 2 - \aleph, & \text{if } \aleph \in [0, 2], \\ 2, & \text{if } \aleph \in \mathbf{N} - \{1, 2\}; \end{cases}$$

$$\mathcal{Q}(\aleph) = \begin{cases} \aleph, & \text{if } \aleph \in [0, 2], \\ 2, & \text{if } \aleph \in \mathbf{N} - \{1, 2\}, \end{cases}$$

for all $\aleph \in \Xi \cup \Omega$. Now, suppose that $\theta = \frac{1}{2}$, then for all $\varphi > 0$, we discuss the following cases:

Case 1. If $\aleph \in [0, 2]$ and $\varpi \in \mathbf{N} - \{1, 2\}$, then

$$\begin{aligned} \mathbb{H}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) &= \mathbb{H}(2 - \aleph, 2, \theta\varphi) = \frac{\theta\varphi}{\theta\varphi + |2 - \aleph - 2|} \\ &\geq \frac{\varphi}{\varphi + |\aleph - \varpi|} = \mathbb{H}(\aleph, \varpi, \varphi), \\ \mathbb{O}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) &= \mathbb{O}(2 - \aleph, 2, \theta\varphi) = \frac{|2 - \aleph - 2|}{\theta\varphi + |2 - \aleph - 2|} \\ &\leq \frac{|\aleph - \varpi|}{\varphi + |\aleph - \varpi|} = \mathbb{O}(\aleph, \varpi, \varphi), \\ \mathbb{S}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) &= \mathbb{S}(2 - \aleph, 2, \theta\varphi) = \frac{|2 - \aleph - 2|}{\theta\varphi} \\ &\leq \frac{|\aleph - \varpi|}{\varphi} = \mathbb{S}(\aleph, \varpi, \varphi). \end{aligned}$$

Case 2. If $\aleph \in \mathbf{N} - \{1, 2\}$ and $\varpi \in [0, 2]$, then

$$\begin{aligned} \mathbb{H}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) &= \mathbb{H}(2, \varpi, \theta\varphi) = \frac{\theta\varphi}{\theta\varphi + |2 - \varpi|} \\ &\geq \frac{\varphi}{\varphi + |\aleph - \varpi|} = \mathbb{H}(\aleph, \varpi, \varphi), \\ \mathbb{O}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) &= \mathbb{O}(2, \varpi, \theta\varphi) = \frac{|2 - \varpi|}{\theta\varphi + |2 - \varpi|} \\ &\leq \frac{|\aleph - \varpi|}{\varphi + |\aleph - \varpi|} = \mathbb{O}(\aleph, \varpi, \varphi), \end{aligned}$$

$$\begin{aligned}\mathbb{S}(\mathcal{P}(\aleph), \mathcal{Q}(\varpi), \theta\varphi) &= \mathbb{S}(2, \varpi, \theta\varphi) = \frac{|2 - \varpi|}{\theta\varphi} \\ &\leq \frac{|\aleph - \varpi|}{\varphi} = \mathbb{S}(\aleph, \varpi, \varphi).\end{aligned}$$

Therefore, all the conditions are fulfilled by $\mathcal{P}$ and $\mathcal{Q}$. By Theorem 3.10, $\mathcal{P}$ and $\mathcal{Q}$ have a unique common fixed point, i.e., $\aleph = 1$.

Definition 3.12. Let $(\Omega, \Xi, \mathbb{M}, *, \circ)$ be a neutrosophic fuzzy bipolar metric space. A map $\mathcal{P} : \Omega \cup \Xi \rightarrow \Omega \cup \Xi$ is an NFB (neutrosophic fuzzy bipolar)-contraction if we can find $0 < \theta < 1$ satisfying

$$\begin{aligned}\frac{1}{\mathbb{H}(\mathcal{P}\aleph, \mathcal{P}\varpi, \varphi)} - 1 &\leq \theta \left[\frac{1}{\mathbb{H}(\aleph, \varpi, \varphi)} - 1 \right], \quad \mathbb{O}(\mathcal{P}\aleph, \mathcal{P}\varpi, \varphi) \leq \theta \mathbb{O}(\aleph, \varpi, \varphi) \\ \mathbb{S}(\mathcal{P}\aleph, \mathcal{P}\varpi, \varphi) &\leq \theta \mathbb{S}(\aleph, \varpi, \varphi), \text{ for all } \aleph \in \Omega, \varpi \in \Xi \text{ and } \varphi > 0.\end{aligned}$$

Theorem 3.13. Let $(\Omega, \Xi, \mathbb{M}, *, \circ)$ be a complete neutrosophic fuzzy bipolar metric space. Let $\mathcal{P} : \Omega \cup \Xi \rightarrow \Omega \cup \Xi$ be a mapping satisfying:

- (1) $\mathcal{P}(\Omega) \subseteq \Omega$ and $\mathcal{P}(\Xi) \subseteq \Xi$;
- (2) $\mathcal{P}$ is IFB-contraction, for all $\aleph \in \Omega$, $\varpi \in \Xi$, and $\varphi > 0$.

Then, $\mathcal{P}$ has a unique fixed point.

Proof. Let $\aleph_0 \in \Omega$ and $\varpi_0 \in \Xi$ and assume that $\mathcal{P}(\aleph_\alpha) = \aleph_{\alpha+1}$ and $\mathcal{P}(\varpi_\alpha) = \varpi_{\alpha+1}$ for all $\alpha \in \mathbb{N} \cup \{0\}$.

Then, we get $(\aleph_\alpha, \varpi_\alpha)$ as a bisequence on the neutrosophic fuzzy bipolar metric space $(\Omega, \Xi, \mathbb{M}, *, \circ)$. By using contraction for all $\varphi > 0$, we deduce

$$\begin{aligned}\frac{1}{\mathbb{H}(\aleph_\alpha, \varpi_\alpha, \varphi)} - 1 &= \frac{1}{\mathbb{H}(\mathcal{P}\aleph_{\alpha-1}, \mathcal{P}\varpi_{\alpha-1}, \varphi)} - 1 \\ &\leq \theta \left[\frac{1}{\mathbb{H}(\aleph_{\alpha-1}, \varpi_{\alpha-1}, \varphi)} - 1 \right] \\ &= \frac{\theta}{\mathbb{H}(\aleph_{\alpha-1}, \varpi_{\alpha-1}, \varphi)} - \theta,\end{aligned}$$

implies that,

$$\begin{aligned}\frac{1}{\mathbb{H}(\aleph_\alpha, \varpi_\alpha, \varphi)} &\leq \frac{\theta}{\mathbb{H}(\aleph_{\alpha-1}, \varpi_{\alpha-1}, \varphi)} + (1 - \theta) \\ &\leq \frac{\theta^2}{\mathbb{H}(\aleph_{\alpha-2}, \varpi_{\alpha-2}, \varphi)} + \theta(1 - \theta) + (1 - \theta).\end{aligned}$$

Continuing this way, we deduce

$$\frac{1}{\mathbb{H}(\aleph_\alpha, \varpi_\alpha, \varphi)} \leq \frac{\theta^\alpha}{\mathbb{H}(\aleph_0, \varpi_0, \varphi)} + \theta^{\alpha-1}(1 - \theta) + \theta^{\alpha-2}(1 - \theta) + \cdots + \theta(1 - \theta) + (1 - \theta)$$

$$\begin{aligned}
&\leq \frac{\theta^\alpha}{\mathbb{H}(\aleph_0, \varpi_0, \varphi)} + (\theta^{\alpha-1} + \theta^{\alpha-2} + \cdots + \theta + 1)(1 - \theta). \\
&\leq \frac{\theta^\alpha}{\mathbb{H}(\aleph_0, \varpi_0, \varphi)} + (1 - \theta^\alpha).
\end{aligned}$$

We obtain

$$\frac{1}{\frac{\theta^\alpha}{\mathbb{H}(\aleph_0, \varpi_0, \varphi)} + (1 - \theta^\alpha)} \leq \mathbb{H}(\aleph_\alpha, \varpi_\alpha, \varphi),$$

$$\begin{aligned}
\mathbb{O}(\aleph_\alpha, \varpi_\alpha, \varphi) &= \mathbb{O}(\mathcal{P}\aleph_{\alpha-1}, \mathcal{P}\varpi_{\alpha-1}, \varphi) \leq \theta \mathbb{O}(\aleph_{\alpha-1}, \varpi_{\alpha-1}, \varphi) = \mathbb{O}(\mathcal{P}\aleph_{\alpha-2}, \mathcal{P}\varpi_{\alpha-2}, \varphi) \\
&\leq \theta^2 \mathbb{O}(\aleph_{\alpha-2}, \varpi_{\alpha-2}, \varphi) \leq \cdots \leq \theta^\alpha \mathbb{O}(\aleph_0, \varpi_0, \varphi) \\
\mathbb{O}(\aleph_{\alpha+1}, \varpi_\alpha, \varphi) &= \mathbb{O}(\mathcal{P}\aleph_\alpha, \mathcal{P}\varpi_{\alpha-1}, \varphi) \leq \theta \mathbb{O}(\aleph_\alpha, \varpi_{\alpha-1}, \varphi) = \mathbb{O}(\mathcal{P}\aleph_{\alpha-1}, \mathcal{P}\varpi_{\alpha-2}, \varphi) \\
&\leq \theta^2 \mathbb{O}(\aleph_{\alpha-1}, \varpi_{\alpha-2}, \varphi) \leq \cdots \leq \theta^\alpha \mathbb{O}(\aleph_1, \varpi_0, \varphi), \\
\mathbb{S}(\aleph_\alpha, \varpi_\alpha, \varphi) &= \mathbb{S}(\mathcal{P}\aleph_{\alpha-1}, \mathcal{P}\varpi_{\alpha-1}, \varphi) \leq \theta \mathbb{S}(\aleph_{\alpha-1}, \varpi_{\alpha-1}, \varphi) = \mathbb{S}(\mathcal{P}\aleph_{\alpha-2}, \mathcal{P}\varpi_{\alpha-2}, \varphi) \\
&\leq \theta^2 \mathbb{S}(\aleph_{\alpha-2}, \varpi_{\alpha-2}, \varphi) \leq \cdots \leq \theta^\alpha \mathbb{S}(\aleph_0, \varpi_0, \varphi) \\
\mathbb{S}(\aleph_{\alpha+1}, \varpi_\alpha, \varphi) &= \mathbb{S}(\mathcal{P}\aleph_\alpha, \mathcal{P}\varpi_{\alpha-1}, \varphi) \leq \theta \mathbb{S}(\aleph_\alpha, \varpi_{\alpha-1}, \varphi) = \mathbb{S}(\mathcal{P}\aleph_{\alpha-1}, \mathcal{P}\varpi_{\alpha-2}, \varphi) \\
&\leq \theta^2 \mathbb{S}(\aleph_{\alpha-1}, \varpi_{\alpha-2}, \varphi) \leq \cdots \leq \theta^\alpha \mathbb{S}(\aleph_1, \varpi_0, \varphi).
\end{aligned}$$

Let $\alpha < \kappa$, for $\alpha, \kappa \in \mathbf{N}$. Then,

$$\begin{aligned}
\mathbb{H}(\aleph_\alpha, \varpi_\kappa, \varphi) &\geq \mathbb{H}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\kappa, \frac{\varphi}{3}\right) \cdots \geq \mathbb{H}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \\
&\quad \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{H}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \\
&\quad \mathbb{H}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right), \\
\mathbb{O}(\aleph_\alpha, \varpi_\kappa, \varphi) &\leq \mathbb{O}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\kappa, \frac{\varphi}{3}\right) \cdots \leq \mathbb{O}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \\
&\quad \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \cdots * \mathbb{O}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{O}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \\
&\quad \mathbb{O}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right)
\end{aligned}$$

and

$$\begin{aligned}
\mathbb{S}(\aleph_\alpha, \varpi_\kappa, \varphi) &\leq \mathbb{S}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\kappa, \frac{\varphi}{3}\right) \cdots \leq \mathbb{S}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \\
&\quad \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \cdots * \mathbb{S}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{S}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{S}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right).
\end{aligned}$$

Therefore,

$$\mathbb{H}(\aleph_\alpha, \varpi_\kappa, \varphi) \geq \mathbb{H}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) * \mathbb{H}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) * \cdots * \mathbb{H}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) * \mathbb{H}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) *$$

$$\begin{aligned} \mathbb{H}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{k-1}}\right) &\geq \frac{1}{\frac{\theta^\alpha}{\mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3}\right)} + (1 - \theta^\alpha)} * \frac{1}{\frac{\theta^\alpha}{\mathbb{H}\left(\aleph_1, \varpi_0, \frac{\varphi}{3}\right)} + (1 - \theta^\alpha)} * \\ &\quad \frac{1}{\frac{\theta^\alpha}{\mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{k-1}}\right)} + (1 - \theta^{k-1})} * \frac{1}{\frac{\theta^\alpha}{\mathbb{H}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{k-1}}\right)} + (1 - \theta^{k-1})} * \\ &\quad \frac{1}{\frac{\theta^\alpha}{\mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{k-1}}\right)} + (1 - \theta^k)}, \end{aligned}$$

$$\begin{aligned} \mathbb{O}(\aleph_\alpha, \varpi_\kappa, \varphi) &\leq \mathbb{O}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{O}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \dots \circ \mathbb{O}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \\ &\quad \mathbb{O}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \mathbb{O}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right) \circ \leq \theta^\alpha \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3}\right) \circ \\ &\quad \theta^\alpha \mathbb{O}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \dots \theta^{\kappa-1} \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \theta^{\kappa-1} \mathbb{O}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \\ &\quad \theta^\kappa \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \end{aligned}$$

and

$$\begin{aligned} \mathbb{S}(\aleph_\alpha, \varpi_\kappa, \varphi) &\leq \mathbb{S}\left(\aleph_\alpha, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \mathbb{S}\left(\aleph_{\alpha+1}, \varpi_\alpha, \frac{\varphi}{3}\right) \circ \dots \circ \mathbb{S}\left(\aleph_{\kappa-1}, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \\ &\quad \mathbb{S}\left(\aleph_\kappa, \varpi_{\kappa-1}, \frac{\varphi}{3^{\kappa-1}}\right) \circ \mathbb{S}\left(\aleph_\kappa, \varpi_\kappa, \frac{\varphi}{3^{\kappa-1}}\right) \circ \leq \theta^\alpha \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3}\right) \circ \theta^\alpha \mathbb{S}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \dots \\ &\quad \theta^{\kappa-1} \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \theta^{\kappa-1} \mathbb{S}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \theta^\kappa \mathbb{S}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right), \end{aligned}$$

implies that,

$$\begin{aligned} \mathbb{H}(\aleph_k, \varpi_k, \varphi) &\geq \frac{1}{\frac{\theta^\alpha}{\mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3}\right)} + (1 - \theta^\alpha)} * \frac{1}{\frac{\theta^\alpha}{\mathbb{H}\left(\aleph_1, \varpi_0, \frac{\varphi}{3}\right)} + (1 - \theta^\alpha)} * \\ &\quad \frac{1}{\frac{\theta^{\kappa-1}}{\mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{k-1}}\right)} + (1 - \theta^{k-1})} * \frac{1}{\frac{\theta^{\kappa-1}}{\mathbb{H}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{k-1}}\right)} + (1 - \theta^{k-1})} * \\ &\quad \frac{1}{\frac{\theta^\kappa}{\mathbb{H}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{k-1}}\right)} + (1 - \theta^k)}, \end{aligned}$$

$$\begin{aligned} \mathbb{O}(\aleph_k, \varpi_k, \varphi) &= \theta^\alpha \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3}\right) \circ \theta^\alpha \mathbb{O}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \dots \\ &\quad \theta^{\kappa-1} \mathbb{O}\left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \theta^{\kappa-1} \mathbb{O}\left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}}\right) \circ \end{aligned}$$

$$\theta^\kappa \mathbb{O} \left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}} \right)$$

and

$$\begin{aligned} \mathbb{S}(\aleph_k, \varpi_k, \varphi) &= \theta^\alpha \mathbb{S} \left(\aleph_0, \varpi_0, \frac{\varphi}{3} \right) \circ \theta^\alpha \mathbb{S} \left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}} \right) \circ \dots \\ &\quad \circ \theta^{\kappa-1} \mathbb{S} \left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}} \right) \circ \theta^{\kappa-1} \mathbb{S} \left(\aleph_1, \varpi_0, \frac{\varphi}{3^{\kappa-1}} \right) \circ \theta^\kappa \mathbb{S} \left(\aleph_0, \varpi_0, \frac{\varphi}{3^{\kappa-1}} \right). \end{aligned}$$

As $\alpha, \kappa \rightarrow +\infty$, we deduce

$$\begin{aligned} \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{H}(\aleph_\alpha, \varpi_\kappa, \varphi) &= 1 * 1 * \dots * 1 = 1, \\ \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{O}(\aleph_\alpha, \varpi_\kappa, \varphi) &= 0 \circ 0 \circ \dots \circ 0 = 0 \text{ and} \\ \lim_{\alpha, \kappa \rightarrow +\infty} \mathbb{S}(\aleph_\alpha, \varpi_\kappa, \varphi) &= 0 \circ 0 \circ \dots \circ 0 = 0. \end{aligned}$$

Which implies that the bisequence $(\aleph_\alpha, \varpi_\alpha)$ is a Cauchy bisequence. Since $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is a complete neutrosophic fuzzy bipolar metric space, then $\{\aleph_\alpha\} \rightarrow \aleph$ and $\{\varpi_\alpha\} \rightarrow \varpi$, where $\aleph \in \Omega \cap \Xi$.

$$\begin{aligned} \mathbb{H}(\aleph, \mathcal{P}\aleph, \varphi) &\geq \mathbb{H} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) * \mathbb{H} \left(\aleph_{\alpha+1}, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) * \mathbb{H} \left(\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\ &= \mathbb{H} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) * \mathbb{H} \left(\mathcal{P}\aleph_\alpha, \mathcal{P}\aleph_\alpha, \frac{\varphi}{3} \right) * \mathbb{H} \left(\mathcal{P}\aleph_\alpha, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\ &\geq \mathbb{H} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) * \frac{1}{\frac{\theta^{\alpha+1}}{\mathbb{H} \left(\aleph_0, \varpi_0, \frac{\varphi}{3} \right)} + (1 - \theta^{\alpha+1})} * \mathbb{H} \left(\mathcal{P}\aleph_\alpha, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\ &\rightarrow 1 * 1 * 1 = 1 \text{ as } \alpha \rightarrow \infty, \end{aligned}$$

$$\begin{aligned} \mathbb{O}(\aleph, \mathcal{P}\aleph, \varphi) &\leq \mathbb{O} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{O} \left(\aleph_{\alpha+1}, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{O} \left(\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\ &= \mathbb{O} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{O} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{O} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\ &\leq \mathbb{O} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \theta^{\alpha+1} \mathbb{O} \left(\aleph_0, \varpi_0, \frac{\varphi}{3} \right) \circ \mathbb{O} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \end{aligned}$$

and

$$\begin{aligned} \mathbb{S}(\aleph, \mathcal{P}\aleph, \varphi) &\leq \mathbb{S} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\aleph_{\alpha+1}, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\ &= \mathbb{S} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right) \\ &\leq \mathbb{S} \left(\aleph, \aleph_{\alpha+1}, \frac{\varphi}{3} \right) \circ \theta^{\alpha+1} \mathbb{S} \left(\aleph_0, \varpi_0, \frac{\varphi}{3} \right) \circ \mathbb{S} \left(\mathcal{P}\aleph_{\alpha+1}, \mathcal{P}\aleph, \frac{\varphi}{3} \right). \end{aligned}$$

Hence, $\mathcal{P}\aleph = \aleph$. Let $\mathcal{P}\varpi = \varpi$ for some $\varpi \in \Omega$, then

$$\frac{1}{\mathbb{H}(\aleph, \omega, \varphi)} - 1 \leq \frac{1}{\mathbb{H}(\mathcal{P}\aleph, \mathcal{P}\omega, \varphi)} - 1 \leq \theta \left[\frac{1}{\mathbb{H}(\aleph, \omega, \varphi)} - 1 \right] < \frac{1}{\mathbb{H}(\aleph, \omega, \varphi)} - 1,$$

$$\mathbb{O}(\aleph, \omega, \varphi) = \mathbb{O}(\mathcal{P}\aleph, \mathcal{P}\omega, \varphi) \leq \theta \mathbb{O}(\aleph, \omega, \varphi) < \mathbb{O}(\aleph, \omega, \varphi)$$

and

$$\mathbb{S}(\aleph, \omega, \varphi) = \mathbb{S}(\mathcal{P}\aleph, \mathcal{P}\omega, \varphi) \leq \theta \mathbb{S}(\aleph, \omega, \varphi) < \mathbb{S}(\aleph, \omega, \varphi),$$

this is a contraction as well. Therefore, $\mathbb{H}(\aleph, \varpi, \varphi) = 1$, $\mathbb{O}(\aleph, \varpi, \varphi) = 0$, $\mathbb{S}(\aleph, \varpi, \varphi) = 0$. Hence, $\aleph = \varpi$. $\square$

Example 3.14. Let $\Omega = [0, 1]$ and $\Xi = \{0\} \cup \mathbf{N} - \{1\}$. Define $\mathbb{M} : \Omega \times \Xi \times (0, +\infty) \rightarrow [0, 1]$ as

$$\mathbb{H}(\aleph, \varpi, \varphi) = \frac{\varphi}{\varphi + |\aleph - \varpi|}, \quad \mathbb{O}(\aleph, \varpi, \varphi) = \frac{|\aleph - \varpi|}{\varphi + |\aleph - \varpi|}, \quad \mathbb{S}(\aleph, \varpi, \varphi) = \frac{|\aleph - \varpi|}{\varphi}.$$

Then, $(\Omega, \Xi, \mathbb{M}, *, \circ)$ is a complete neutrosophic fuzzy bipolar metric space with CTN $\nu * \aleph = \nu \aleph$ and CTCN $\nu \circ \aleph = \max\{\nu, \aleph\}$.

Proof. Define,

$$\mathcal{P} : (\Omega, \Xi, \mathbb{M}, *, \circ) \rightrightarrows (\Omega, \Xi, \mathbb{M}, *, \circ)$$

by

$$\mathcal{P}(\aleph) = \begin{cases} \frac{1 - 3^{-\aleph}}{5}, & \text{if } \aleph \in [0, 1] \\ 0, & \text{if } \aleph \in \mathbf{N} - \{1\}, \end{cases}$$

for all $\aleph \in \Omega \cup \Xi$ and take $\theta \in [\frac{1}{2}, 1)$, then

$$\begin{aligned} \mathbb{H}(\mathcal{P}\aleph, \mathcal{P}\varpi, \theta\varphi) &= \mathbb{H}\left(\frac{1 - 3^{-\aleph}}{5}, \frac{1 - 3^{-\varpi}}{5}, \theta\varphi\right) = \frac{\theta\varphi}{\theta\varphi + \left|\frac{1 - 3^{-\aleph}}{5} - \frac{1 - 3^{-\varpi}}{5}\right|} \\ &= \frac{\theta\varphi}{\theta\varphi + \left|\frac{3^{-\aleph} - 3^{-\varpi}}{5}\right|} \geq \frac{\theta\varphi}{\theta\varphi + \frac{|\aleph - \varpi|}{5}} \\ &= \frac{5\theta\varphi}{5\theta\varphi + |\aleph - \varpi|} \geq \frac{\varphi}{\varphi + |\aleph - \varpi|} = \mathbb{H}(\aleph, \varpi, \varphi), \end{aligned}$$

$$\mathbb{O}(\mathcal{P}\aleph, \mathcal{P}\varpi, \theta\varphi) = \mathbb{O}\left(\frac{1 - 3^{-\aleph}}{5}, \frac{1 - 3^{-\varpi}}{5}, \theta\varphi\right) = \frac{\left|\frac{1 - 3^{-\aleph}}{5} - \frac{1 - 3^{-\varpi}}{5}\right|}{\theta\varphi + \left|\frac{1 - 3^{-\aleph}}{5} - \frac{1 - 3^{-\varpi}}{5}\right|}$$

$$\begin{aligned}
&= \frac{\frac{|3^{-\aleph} - 3^{-\varpi}|}{5}}{\theta\varphi + \frac{|3^{-\aleph} - 3^{-\varpi}|}{5}} \leq \frac{|3^{-\aleph} - 3^{-\varpi}|}{5\theta\varphi + |3^{-\aleph} - 3^{-\varpi}|} \\
&\leq \frac{|\aleph - \varpi|}{5\theta\varphi + |\aleph - \varpi|} \leq \frac{|\aleph - \varpi|}{\theta\varphi + |\aleph - \varpi|} = \mathbb{O}(\aleph, \varpi, \varphi)
\end{aligned}$$

and

$$\begin{aligned}
\mathbb{S}(\mathcal{P}\aleph, \mathcal{P}\varpi, \theta\varphi) &= \mathbb{S}\left(\frac{1 - 3^{-\aleph}}{5}, \frac{1 - 3^{-\varpi}}{5}, \theta\varphi\right) = \frac{\left|\frac{1 - 3^{-\aleph}}{5} - \frac{1 - 3^{-\varpi}}{5}\right|}{\theta\varphi} \\
&= \frac{\frac{|3^{-\aleph} - 3^{-\varpi}|}{5}}{\theta\varphi} \leq \frac{|3^{-\aleph} - 3^{-\varpi}|}{5\theta\varphi} \leq \frac{|\aleph - \varpi|}{5\theta\varphi} \\
&\leq \frac{|\aleph - \varpi|}{\theta\varphi} = \mathbb{S}(\aleph, \varpi, \varphi).
\end{aligned}$$

Therefore, the conditions of Theorem 3.13 are fulfilled, and 0 is the only fixed point for $\mathcal{P}$. $\square$

4. APPLICATION

Let $\Omega = \mathcal{C}([c, a], [0, +\infty))$ be the set of all continuous functions defined on $[c, a]$ with values in the interval $[0, +\infty)$, and $\Xi = \mathcal{C}([c, a], (-\infty, 0])$ be the set of all continuous functions defined on $[c, a]$ with values in the interval $(-\infty, 0]$.

Theorem 4.1. *Suppose the integral equation:*

$$\aleph(\chi) = \varpi(\chi) + \delta \int_a^c \mathcal{R}(\chi, \rho) \aleph(\chi) d\rho \quad \text{for } \chi, \rho \in [c, a], \quad (2)$$

where $\delta > 0$, $\varpi(\rho)$ is a fuzzy function of $\rho : \rho \in [c, a]$ and $\mathcal{R} : \mathcal{C}([c, a] \times \mathbf{R}) \rightarrow \mathbf{R}^+$.

Define, $\mathbb{H}$, $\mathbb{O}$ and $\mathbb{S}$ by

$$\mathbb{H}(\aleph(\chi), \varpi(\chi), \varphi) = \begin{cases} \frac{\varphi}{\varphi + \max_{\chi \in [c, a]} |\aleph(\chi) - \varpi(\chi)|} & \text{if } \varphi > 0 \\ 0 & \text{if } \varphi = 0; \end{cases}$$

$$\mathbb{O}(\aleph(\chi), \varpi(\chi), \varphi) = \begin{cases} \frac{\max_{\chi \in [c, a]} |\aleph(\chi) - \varpi(\chi)|}{\varphi + \max_{\chi \in [c, a]} |\aleph(\chi) - \varpi(\chi)|} & \text{if } \varphi > 0 \\ 0 & \text{if } \varphi = 0 \end{cases}$$

and

$$\mathbb{S}(\aleph(\chi), \varpi(\chi), \varphi) = \begin{cases} \frac{\max_{\chi \in [c, a]} |\aleph(\chi) - \varpi(\chi)|}{\varphi} & \text{if } \varphi > 0, \\ 0 & \text{if } \varphi = 0 \end{cases}$$

with CTN and CTCN defined by $\nu * \aleph = \nu \cdot \aleph$ and $\nu \circ \aleph = \max(\nu, \aleph)$. Then $(\Xi, \Omega, \mathbb{M}, *, \circ)$ is a complete neutrosophic fuzzy bipolar metric space.

Suppose that $|\mathcal{R}(\chi, \rho)\aleph(\chi) - \mathcal{R}(\chi, \rho)\varpi(\chi)| \leq |\aleph(\chi) - \varpi(\chi)|$ for $\aleph \in \Omega$, $\varpi \in \Xi$, $\theta \in (0, 1)$ and all $\chi, \rho \in [c, a]$. Also, let $\mathcal{R}(\chi, \rho)(\delta \int_c^a d\rho) \leq \theta < 1$.

Then, the integral Equation (2) has a unique solution.

Proof. Define $\mathcal{P} : (\Xi, \Omega, \mathbb{H}, \mathbb{O}, *, \circ) \rightarrow (\Xi, \Omega, \mathbb{H}, \mathbb{O}, *, \circ)$ by

$$\mathcal{P}\aleph(\chi) = \varpi(\chi) + \delta \int_c^a \mathcal{R}(\chi, \rho)\aleph(\chi)d\rho, \text{ for all } \chi, \rho \in [c, a].$$

Now, for all $\aleph, \varpi \in \Omega \cup \Xi$, we deduce

$$\begin{aligned} \mathbb{H}(\mathcal{P}\aleph(\chi), \mathcal{P}\varpi(\chi), \theta\varphi) &= \frac{\theta\varphi}{\theta\varphi + \max_{\chi \in [c, a]} |\mathcal{P}\aleph(\chi) - \mathcal{P}\varpi(\chi)|} \\ &= \frac{\theta\varphi}{\theta\varphi + \max_{\chi \in [c, a]} [|\varpi(\chi) + \delta \int_c^a \mathcal{R}(\chi, \rho)\aleph(\chi)d\rho - \varpi(\chi) - \delta \int_c^a \mathcal{R}(\chi, \rho)\varpi(\chi)d\rho|]} \\ &= \frac{\theta\varphi}{\theta\varphi + \max_{\chi \in [c, a]} [|\delta \int_c^a \mathcal{R}(\chi, \rho)\aleph(\chi)d\rho - \delta \int_c^a \mathcal{R}(\chi, \rho)\varpi(\chi)d\rho|]} \\ &= \frac{\theta\varphi}{\theta\varphi + \max_{\chi \in [c, a]} [|\mathcal{R}(\chi, \rho)\aleph(\chi) - \mathcal{R}(\chi, \rho)\varpi(\chi)| (\delta \int_c^a d\rho)]} \\ &\geq \frac{\varphi}{\varphi + \max_{\chi \in [c, a]} [|\aleph(\chi) - \varpi(\chi)|]} \\ &\geq \mathbb{H}(\aleph(\chi), \varpi(\chi), \varphi), \end{aligned}$$

$$\begin{aligned} \mathbb{O}(\mathcal{P}\aleph(\chi), \mathcal{P}\varpi(\chi), \theta\varphi) &= \frac{\max_{\chi \in [c, a]} |\mathcal{P}\aleph(\chi) - \mathcal{P}\varpi(\chi)|}{\theta\varphi + \max_{\chi \in [c, a]} |\mathcal{P}\aleph(\chi) - \mathcal{P}\varpi(\chi)|} \\ &= \frac{\max_{\chi \in [c, a]} [|\varpi(\chi) + \delta \int_c^a \mathcal{R}(\chi, \rho)\aleph(\chi)d\rho - \varpi(\chi) - \delta \int_c^a \mathcal{R}(\chi, \rho)\varpi(\chi)d\rho|]}{\theta\varphi + \max_{\chi \in [c, a]} [|\varpi(\chi) + \delta \int_c^a \mathcal{R}(\chi, \rho)\aleph(\chi)d\rho - \varpi(\chi) - \delta \int_c^a \mathcal{R}(\chi, \rho)\varpi(\chi)d\rho|]} \\ &= \frac{\max_{\chi \in [c, a]} [|\delta \int_c^a \mathcal{R}(\chi, \rho)\aleph(\chi)d\rho - \delta \int_c^a \mathcal{R}(\chi, \rho)\varpi(\chi)d\rho|]}{\theta\varphi + \max_{\chi \in [c, a]} [|\delta \int_c^a \mathcal{R}(\chi, \rho)\aleph(\chi)d\rho - \delta \int_c^a \mathcal{R}(\chi, \rho)\varpi(\chi)d\rho|]} \\ &= \frac{\max_{\chi \in [c, a]} [|\mathcal{R}(\chi, \rho)\aleph(\chi) - \mathcal{R}(\chi, \rho)\varpi(\chi)| (\delta \int_c^a d\rho)]}{\theta\varphi + \max_{\chi \in [c, a]} [|\mathcal{R}(\chi, \rho)\aleph(\chi) - \mathcal{R}(\chi, \rho)\varpi(\chi)| (\delta \int_c^a d\rho)]} \\ &\geq \frac{\max_{\chi \in [c, a]} [|\aleph(\chi) - \varpi(\chi)|]}{\varphi + \max_{\chi \in [c, a]} [|\aleph(\chi) - \varpi(\chi)|]} \geq \mathbb{O}(\aleph(\chi), \varpi(\chi), \varphi) \end{aligned}$$

and

$$\mathbb{S}(\mathcal{P}\aleph(\chi), \mathcal{P}\varpi(\chi), \theta\varphi) = \frac{\max_{\chi \in [c, a]} |\mathcal{P}\aleph(\chi) - \mathcal{P}\varpi(\chi)|}{\theta\varphi}$$

$$\begin{aligned}
&= \frac{\max_{\chi \in [c, a]} \left[|\varpi(\chi) + \delta \int_c^a \mathcal{R}(\chi, \rho) \aleph(\chi) d\rho - \varpi(\chi) - \delta \int_c^a \mathcal{R}(\chi, \rho) \varpi(\chi) d\rho| \right]}{\theta \varphi} \\
&= \frac{\max_{\chi \in [c, a]} \left[|\delta \int_c^a \mathcal{R}(\chi, \rho) \aleph(\chi) d\rho - \delta \int_c^a \mathcal{R}(\chi, \rho) \varpi(\chi) d\rho| \right]}{\theta \varphi} \\
&= \frac{\max_{\chi \in [c, a]} \left[|\mathcal{R}(\chi, \rho) \aleph(\chi) - \mathcal{R}(\chi, \rho) \varpi(\chi)| \quad (\delta \int_c^a d\rho) \right]}{\theta \varphi} \\
&\geq \frac{\max_{\chi \in [c, a]} \left[|\aleph(\chi) - \varpi(\chi)| \right]}{\varphi} \geq \mathbb{S}(\aleph(\chi), \varpi(\chi), \varphi).
\end{aligned}$$

Therefore, the conditions of Theorem 3.9 are fulfilled and operator $\mathcal{P}$ has a unique fixed point. $\square$

Example 4.2. Assume the following integral equation:

$$\aleph(\chi) = |\cos \chi| + \frac{1}{7} \int_0^1 \rho \aleph(\rho) d\rho, \quad \text{for all } \rho \in [0, 1].$$

Then, it has a unique solution in $\Omega \cup \Xi$.

Proof. Let $\mathcal{P} : (\Omega, \Xi, \mathbb{M}, *, \circ) \rightrightarrows (\Omega, \Xi, \mathbb{M}, *, \circ)$ be defined by

$$\mathcal{P}\aleph(\chi) = |\sin \chi| + \frac{1}{7} \int_0^1 \rho \aleph(\rho) d\rho,$$

and set $\mathcal{R}(\chi, \rho) \aleph(\chi) = \frac{1}{7} \rho \aleph(\rho)$ and $\mathcal{R}(\chi, \rho) \varpi(\chi) = \frac{1}{7} \rho \varpi(\rho)$, where $\aleph, \varpi \in \Omega \cup \Xi$ and for all $\chi, \rho \in [0, 1]$. Then, we have

$$|\mathcal{R}(\chi, \rho) \aleph(\chi) - \mathcal{R}(\chi, \rho) \varpi(\chi)| = \left| \frac{1}{7} \rho \aleph(\rho) - \frac{1}{7} \rho \varpi(\rho) \right| = \frac{1}{7} \rho |\aleph(\rho) - \varpi(\rho)| \leq |\aleph(\rho) - \varpi(\rho)|,$$

and

$$\frac{1}{7} \int_0^1 \rho d\rho = \frac{1}{7} \left(\frac{(1)^2}{2} - \frac{(0)^2}{2} \right) = \frac{1}{7} = \theta < 1, \quad \text{where } \delta = \frac{1}{7}.$$

Therefore, the conditions of Theorem 3.9 are fulfilled. Hence, $\mathcal{P}$ has a unique solution in $\Omega \cup \Xi$. $\square$

CONCLUSION

This study introduces and explores a novel mathematical structure called the neutrosophic fuzzy bipolar metric space, which extends and generalizes existing fuzzy and bipolar metric space frameworks and formulate new definitions and properties of neutrosophic fuzzy bipolar metric spaces. Establish several fixed point theorems using covariant maps and neutrosophic fuzzy bipolar-contractions. Provide proofs and examples to validate the theoretical findings. Present an application to demonstrate the existence and uniqueness of solutions to certain differential equations within the developed framework. The results unify and generalize many existing fixed point results in fuzzy and neutrosophic metric spaces, making a significant contribution to the fixed point theory in uncertain environments.

FUTURE SCOPE

The present work opens several avenues for future research. One promising direction is the extension to multivalued or set-valued mappings, where fixed point results can be investigated under the framework of neutrosophic fuzzy bipolar metric spaces. Another significant area involves the application of this theory to real-world problems in fields such as engineering, decision-making, and artificial intelligence, especially where uncertainty and bipolar behavior are inherent. Additionally, the current structure can be generalized to other types of metric spaces, including neutrosophic fuzzy bipolar b-metric spaces, cone metric spaces, and probabilistic metric spaces, to broaden the applicability of the theory. From a computational perspective, there is scope for designing algorithmic approaches to numerically compute fixed points within neutrosophic bipolar contexts. Finally, future work could focus on the exploration of dynamic systems, particularly analyzing the stability and convergence behavior of iterative processes within such enriched metric frameworks.

ABBREVIATIONS

FMS-Fuzzy metric space
 FP-Fixed point
 IFS-Intuitionistic fuzzy sets
 IFMS-Intuitionistic fuzzy metric space
 NS-Neutrosophic set
 NMS-Neutrosophic metric space
 MIFMS-modified intuitionistic fuzzy metric space

REFERENCES

- [1] L. A. Zadeh, "Fuzzy sets," *Information and Control*, vol. 8, pp. 338–353, 1965. [https://doi.org/10.1016/S0019-9958\(65\)90241-X](https://doi.org/10.1016/S0019-9958(65)90241-X).
- [2] O. Kaleva and S. Seikkala, "On fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 12, pp. 215–229, 1984.
- [3] B. Schweizer and A. Sklar, "Statistical metric spaces," *Pacific Journal of Mathematics*, vol. 10, no. 1, pp. 313–334, 1960.
- [4] I. Kramosil and J. Michálek, "Fuzzy metrics and statistical metric spaces," *Kybernetika*, vol. 11, no. 5, pp. 336–344, 1975. <http://eudml.org/doc/28711>.
- [5] M. Grabiec, "Fixed points in fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 27, pp. 385–389, 1988.
- [6] J. H. Park, "Intuitionistic fuzzy metric spaces," *Chaos, Solitons & Fractals*, vol. 22, no. 5, pp. 1039–1046, 2004. <https://doi.org/10.1016/j.chaos.2004.02.051>.
- [7] A. George and P. Veeramani, "On some results in fuzzy metric spaces," *Fuzzy Sets and Systems*, vol. 64, pp. 395–399, 1994. <https://www.scirp.org/reference/referencespapers?referenceid=157638>.

- [8] S. Cho and J. Jung, "On common fixed point theorems in fuzzy metric spaces," *International Mathematical Forum*, vol. 1, pp. 1441–1451, 2006. <https://www.m-hikari.com/imf-password/29-32-2006/choIMF29-32-2006.pdf>.
- [9] V. Gregori and A. Sapena, "On fixed-point theorems in fuzzy metric spaces," *Fuzzy sets and systems*, vol. 125, no. 2, pp. 245–252, 2002. [https://doi.org/10.1016/S0165-0114\(00\)00088-9](https://doi.org/10.1016/S0165-0114(00)00088-9).
- [10] V. Gupta, A. Kanwar, and R. Shukla, "Topological properties and fixed point results for modified intuitionistic generalized fuzzy metric spaces," *International Journal of Analysis and Applications*, vol. 23, pp. 1–12, 2025. <https://doi.org/10.28924/2291-8639-23-2025-102>.
- [11] V. Gupta, A. Gondhi, and R. Shukla, "Fixed point results in modified intuitionistic fuzzy soft metric spaces with application," *Mathematics*, vol. 12, no. 8, pp. Article ID:1154, 1–11, 2024.
- [12] V. Gupta and A. Gondhi, "Fixed points of weakly compatible maps on modified intuitionistic fuzzy soft metric spaces," *International Journal of Systems Assurance Engineering and Management*, vol. 13, no. 3, pp. 1232–1238, 2022.
- [13] V. Gupta, A. Saini, and A. Gondhi, "On modified intuitionistic fuzzy soft metric spaces and application," *Miskolc Mathematical Notes*, vol. 25, no. 1, pp. 255–269, 2024. <https://doi.org/10.18514/MMN.2024.4323>.
- [14] A. Mutlu and U. Gürdal, "Bipolar metric spaces and some fixed point theorems," *Journal of nonlinear science and applications*, vol. 9, no. 9, pp. 5362–5373, 2016. <http://dx.doi.org/10.22436/jnsa.009.09.05>.
- [15] K. Roy and M. Saha, "Generalized contractions and fixed point theorems over bipolar cone tvs b-metric spaces with an application to homotopy theory.," *Matematički Vesnik*, vol. 72, no. 4, 2020. <http://elib.mi.sanu.ac.rs/files/journals/mv/281/mvn281p281-296.pdf>.
- [16] Y. U. Gaba, M. Aphane, and H. Aydi, " α , bk-contractions in bipolar metric spaces," *Journal of Mathematics*, vol. 2021, pp. 1–6, 2021. <https://doi.org/10.1155/2021/5562651>.
- [17] U. Gürdal, A. Mutlu, and K. Özkan, "Fixed point results for α - ψ -contractive mappings in bipolar metric spaces," *J. Inequal. Spec. Funct.*, vol. 11, no. 1, pp. 64–75, 2020. www.ilirias.com/jiasf.
- [18] G. Kishore, R. P. Agarwal, B. Srinuvasa Rao, and R. Srinivasa Rao, "Caristi type cyclic contraction and common fixed point theorems in bipolar metric spaces with applications," *Fixed point theory and Applications*, vol. 2018, no. 1, p. 21, 2018. <https://doi.org/10.1186/s13663-018-0646-z>.
- [19] G. N. V. Kishore, D. R. Prasad, B. S. Rao, and V. S. Baghavan, "Some applications via common coupled fixed point theorems in bipolar metric spaces," *Journal of Critical Reviews*, vol. 7, pp. 601–607, 2020. <https://doi.org/10.31838/jcr.07.02.110>.
- [20] G. Kishore, K. Rao, H. Isik, B. Srinuvasa Rao, and A. Sombabu, "Covarian mappings and coupled fixed point results in bipolar metric spaces," *International Journal of Nonlinear Analysis and Applications*, vol. 12, no. 1, pp. 1–15, 2021. <https://doi.org/10.22075/ijnaa.2021.4650>.
- [21] G. Kishore, K. Rao, A. Sombabu, and R. Rao, "Related results to hybrid pair of mappings and applications in bipolar metric spaces," *Journal of Mathematics*, vol. 2019, pp. 1–7, 2019. <https://doi.org/10.1155/2019/8485412>.
- [22] G. Mani, A. J. Gnanaprakasam, N. Kausar, M. Munir, S. Khan, and E. Ozbilge, "Solving an integral equation via intuitionistic fuzzy bipolar metric spaces," *Decision Making: Applications in Management and Engineering*, vol. 6, no. 2, pp. 536–556, 2023. <https://doi.org/10.31181/dmame622023624>.
- [23] A. Mutlu, K. Özkan, and U. Gürdal, "Locally and weakly contractive principle in bipolar metric spaces," *TWMS Journal of Applied and Engineering Mathematics*, vol. 10, no. 2, pp. 379–388, 2020. <https://jaem.isikun.edu.tr/web/images/articles/vol.10.no.2/11.pdf>.
- [24] B. S. Rao, G. Kishore, and G. K. Kumar, "Geraghty type contraction and common coupled fixed point theorems in bipolar metric spaces with applications to homotopy," *International*

- Journal of Mathematics Trends and Technology-IJMTT*, vol. 63, pp. 25–34, 2018. <https://doi.org/10.14445/22315373/IJMTT-V63P504>.
- [25] K. Roy and M. Saha, “Sequential bipolar metric space and well-posedness of fixed point problems,” *International Journal of Nonlinear Analysis and Applications*, vol. 12, no. 2, pp. 387–398, 2021. <https://doi.org/10.22075/ijnaa.2020.20397.2150>.
 - [26] K. Roy, M. Saha, R. George, L. Guran, and Z. D. Mitrović, “Some covariant and contravariant fixed point theorems over bipolar metric spaces and applications,” *Filomat*, vol. 36, no. 5, pp. 1755–1767, 2022. <https://doi.org/10.2298/FIL2205755R>.
 - [27] M. Kirisci and N. Simsek, “Neutrosophic metric spaces and fixed point results,” in *Conference Proceedings of Science and Technology*, vol. 2, pp. 64–67, 2019. <https://doi.org/10.48550/arXiv.1907.00798>.
 - [28] V. Gupta, N. Mani, and S. Kapoor, “Contraction theorems in neutrosophic soft metric space,” in *American Institute of Physics Conference Series*, vol. 2735, p. Article ID: 040019, 2023. <https://doi.org/10.1063/5.0140216>.
 - [29] N. Garg and V. Gupta, “Fixed points for compatible maps of type α , β and r-weakly commuting mappings with application in decision making,” *Modeling Earth Systems and Environment*, vol. 11, no. 2, pp. 1–20, 2025. <https://doi.org/10.1007/s40808-024-02225-7>.
 - [30] V. Gupta, N. Garg, and R. Shukla, “Subsequential continuity in neutrosophic metric space with applications,” *Computation*, vol. 13, no. 4, pp. 1–21, 2025. <https://doi.org/10.3390/computation13040087>.
 - [31] S. Sowndrarajan, M. Jeyaraman, and F. Smarandache, *Fixed point results for contraction theorems in neutrosophic metric spaces*, vol. 36. Infinite Study, 2020. https://digitalrepository.unm.edu/nss_journal/vol36/iss1/23.