

ANTIMAGIC LABELING OF GENERALIZED SAUSAGE GRAPHS

OUDONE PHANALASY

School of Mathematical and Physical Sciences
University of Newcastle, Australia

and

Department of Mathematics
National University of Laos, Laos

oudone.phanalasy@gmail.com

Abstract. An *antimagic labeling* of a graph with q edges is a bijection from the set of edges to the set of positive integers $\{1, 2, \dots, q\}$ such that all vertex weights are pairwise distinct, where the *vertex weight* of a vertex is the sum of the labels of all the edges incident with that vertex. A graph is *antimagic* if it has an antimagic labeling. In this paper we construct antimagic labeling for the family of generalized sausage graphs.

Key words: Antimagic labeling, generalized sausage graph.

Abstrak. Sebuah *pelabelan anti-ajaib* dari sebuah graf dengan q sisi adalah sebuah bijeksi dari himpunan sisi-sisi pada himpunan bilangan bulat positif $\{1, 2, \dots, q\}$ sedemikian sehingga semua bobot simpul berbeda per-pasangan, dimana *bobot simpul* dari sebuah simpul adalah jumlah dari label semua sisi yang bersesuaian dengan simpul. Sebuah graf adalah *anti-ajaib* jika dia mempunyai sebuah pelabelan anti-ajaib. Dalam paper ini akan dikonstruksi pelabelan anti-ajaib untuk keluarga graf sosis tergeneralisasi.

Kata kunci: Pelabelan anti-ajaib, graf sosis tergeneralisasi

1. INTRODUCTION

All graphs in this paper are finite, simple, undirected and connected, unless stated otherwise. In 1990, Hartsfield and Ringel [5] introduced the concept of an antimagic labeling of graph, that is, a vertex antimagic edge labeling. An *antimagic*

2000 *Mathematics Subject Classification:* 05C78

Received: 24-12-2013, revised: 13-03-2014, accepted: 14-03-2014.

labeling of a graph $G = (V, E)$ is a bijection $f : E \rightarrow \{1, 2, \dots, |E|\}$ such that all vertex weights are pairwise distinct, where the weight of a vertex v of G , $wt(v)$, is the sum of the labels of all edges incident with the vertex v . A graph G is said to be *antimagic* if it has an antimagic labeling.


Hartsfield and Ringel [5] showed that P_n, S_n, C_n, K_n, W_n , and $K_{2,n}$, for $n \geq 3$, are antimagic. They also conjectured that every connected graph, except K_2 , is antimagic. Subsequently, several families of graphs have been proved to be antimagic, for example, see [1, 2, 3, 10]. Many other results concerning antimagic graphs are catalogued in [4]. Most recently, new families of antimagic graphs have been discovered by Phanalasy *et al.* [7], Miller *et al.* [6] and Rylands *et al.* [9]. However, the conjecture still remains open.

In the previous papers ([7, 9] for example), the results concerned regular and non regular graphs. Here we are extending the method to cover a class of almost regular graphs. We introduce a new family of graphs, called generalized sausage graphs, and we construct antimagic labeling for such family of graphs. The definition of this family of graphs is stated in Section 2.

Hereafter an edge labeling l of a graph G will be described by an array L (not necessary rectangular), where all edge labels incident with a vertex are written in the same row. Since we are dealing with graphs, each label must occur exactly in two different rows.

2. MAIN RESULTS

We first define a new family of graphs. Let G be a k -regular graph with p vertices and q edges. The *generalized sausage graph*, denoted by $S(G, m)$, is the graph obtained from the Cartesian product graph $G \times P_m$, $m \geq 1$ ($G \times P_1 = G$), by joining each vertex of each end of the $G \times P_m$ to a further vertex with an edge; and the two new vertices called *apexes*. In particular, when $m = 1$, each vertex of the graph G joins to two vertices with two edges. The *mixed generalized sausage graph*, denoted by $MS(G, m)$, is the graph obtained from the generalized sausage graph $S(G, m)$, $m \geq 3$, by joining each vertex of each copy of the $\lceil \frac{m}{2} \rceil$ copies of G on the left hand side to the left hand side apex, except the nearest copy to the apex, similarly, for the right hand side apex. The *complete mixed generalized sausage graph*, denoted by $CMS(G, m)$ is the graph obtained from the generalized sausage graph by joining each vertex of each copy of G , except the two nearest copies of G to the apexes, to each apex with an edge, and each corresponding pair of vertices of the two nearest copies of G to the apexes with an edge. The complete mixed generalized sausage graph $CMS^-(G, m)$ is the graph obtained from $CMS(G, m)$ by deleting the edge connecting each corresponding pair of vertices of the two nearest copies of G to the apexes. For an example of the graph $CMS(G, m)$, see Figure 1. Let A, B and C be the sets of the dark dashed edges, tiny dashed edges and dark edges of the graph in Figure 1, respectively. Then the graph $CMS^-(C_3, 5)$ is the


 FIGURE 1. Complete mixed generalized sausage graph $CMS(C_3, 5)$

graph in Figure 1 without A , while the graphs $MS(C_3, 5)$ and $S(C_3, 5)$ are the graphs in Figure 1 without $A \cup B$ and without $A \cup B \cup C$, respectively.

Let G be any (connected or disconnected) k -regular graph with p vertices and q edges. We first choose any labeling of G , that is, label the edges of G by allocating integers $1, 2, \dots, q$ randomly. Then calculate the weights of the vertices and order the vertices so that $wt(v_i) \leq wt(v_{i+1})$, $1 \leq i \leq p - 1$. This ordering results in an array of edge labels of G . It is considered as the original labeling and will be applied throughout the paper to produce antimagic labelings for graphs in the family of generalized sausage graphs.

Denote by T^t the transpose of the array T . Let $T = (1 \ 2 \ \dots \ p - 1 \ p)^t$. We define the *reverse* of the array T as $T^\uparrow = (p \ p - 1 \ \dots \ 2 \ 1)^t$ and

Theorem 2.1. *Let $G \neq nK_1$, $n \geq 1$, be any connected or disconnected k -regular graph. Then the generalized sausage graph $S(G, m)$, $m \geq 1$, is antimagic.*

PROOF. Let L_j , $1 \leq j \leq m$, be the array of edge labels of the j -th copy of the graph G in $S(G, m)$, $m \geq 1$. Let T_l , $1 \leq l \leq m + 1$, be the $(p \times 1)$ -array of the edges e_i , $1 \leq i \leq p$, where e_i are the edges of $S(G, m)$, $m \geq 1$, that do not belong to any copy of G . We construct the array A of edge labels of $S(G, m)$, $m \geq 1$, as follows.

Case 1: $G = K_p$, $p \geq 2$




FIGURE 2. Illustration of antimagic labeling of $S(K_p, m)$, $p \geq 2$, $m \geq 2$ and m even

- (1) Label the edge e_i , $1 \leq i \leq p$, in the row i of the array T_l , $1 \leq l \leq m + 1$, with $i + (l - 1)p$, for $1 \leq l \leq 2$; and $i + (l - 1)p + (l - 2)q$, for $3 \leq l \leq m + 1$;
- (2) Replace the edge labels in the array L_j , $1 \leq j \leq m$, with new labels obtained by adding $(j + 1)p + (j - 1)q$ to each of the original edge labels;
- (3) Form the array A as shown below.
For $m = 1, 2$,

$$\begin{array}{ccc}
 & & T_1^t \\
 & & T_2^t \\
 T_1 & T_2 & L_1 \\
 & & \\
 & & T_1^t \\
 & & T_2^t \\
 T_1 & L_1 & T_3 \\
 T_2 & T_3 & L_2
 \end{array}$$

More generally, for $m \geq 3$,

$$\begin{array}{ccc}
 & & T_1^t \\
 & & T_2^t \\
 T_1 & L_1 & T_3 \\
 T_2 & L_2 & T_4 \\
 \vdots & \vdots & \vdots \\
 T_{m-1} & L_{m-1} & T_{m+1} \\
 T_m & T_{m+1} & L_m
 \end{array}$$

The diagram in Figure 2 illustrates the antimagic labeling used here.

By the construction of the array A , it is clear that the weight of each vertex (row) in the array is less than the weight of the vertex (row) below.

Case 2: $G \neq K_p$, $p \geq 1$

- (1) Replace the edge labels in the array L_j , $1 \leq j \leq m$, with new labels obtained by adding $(j - 1)(p + q)$ to each of the original edge labels;
- (2) Label the edge e_i , $1 \leq i \leq p$, in the row i of the array T_l , $1 \leq l \leq m + 1$, with $i + (l - 1)p + lq$, for $1 \leq l \leq m$; and $i + (l - 1)p + mq$, for $l = m + 1$;
- (3) Form the array A into two cases as shown below (two separate cases).

Subcase 2.1: $m = 1$

$$\begin{array}{ccc}
 L_1 & T_1 & T_2 \\
 & & T_1^t \\
 & & T_2^t
 \end{array}$$

By the construction of the array A , it is clear that the weight of each vertex (row) is less than the weight of the vertex (row) below, except the weight of the last row of the subarray $L_1T_1T_2$ and the weight of the row T_1^t and T_2^t that needs to be verified. This we do in 3 subcases.

Let $wt(r_f)$ be the weight of the row r_f .

Subcase 2.1.1: $G = 2K_2$ ($p = 4, q = 2$ and $k = 1$)

We have $wt(r_p) = wt(T_1^t) = 18$. However, by swapping the edge labels 7 and 10, then all weights of vertices are pairwise distinct.

Subcase 2.1.2: $p = q = 4$ and $k = 2$

It is simple to check that $wt(r_p) = 27, wt(T_1^t) = 26$ and $wt(T_2^t) = 42$.

Subcase 2.1.3: $p, q > 4, k > 1$

Since the largest possible edge labels of the last row in the array L_1 are $q - (k - 1), q - (k - 2), \dots, q - 1$ and q , hence we have $wt(r_p) \leq (3p + k + 2)q - \frac{k(k-1)}{2} < 3p + (k + 2)q < \frac{p(p+1)}{2} + pq = wt(T_1^t)$. It is obvious that $wt(T_1^t) < wt(T_2^t)$.

Subcase 2.2: $m \geq 2$

For $m = 2$,

$$\begin{array}{ccc} L_1 & T_1 & T_2 \\ T_1 & L_2 & T_3 \\ & & T_2^t \\ & & T_3^t \end{array}$$

More generally, for $m \geq 3$,

$$\begin{array}{ccc} L_1 & T_1 & T_2 \\ T_1 & L_2 & T_3 \\ \vdots & \vdots & \vdots \\ T_{m-2} & L_{m-1} & T_m \\ T_{m-1} & L_m & T_{m+1} \\ & & T_m^t \\ & & T_{m+1}^t \end{array}$$

By the construction of the array A , it is clear that the weight of each vertex (row) in the array is less than the weight of the vertex (row) below, except the weights of the last row in the subarray $T_{m-1}L_mT_{m+1}$ and the row T_m^t that need to be verified.

Let $e_{f,g}$ be the edge label in the row f and the column g in the array A . We have the largest possible edge labels in the last row (that is the row r_{mp}) of the array L_m and the row T_m^t as shown below.

$$\begin{array}{llll} r_{mp} : & \dots & (q - 1) + (m - 1)(p + q) & q + (m - 1)(p + q) & (m + 1)p + mq \\ T_m^t : & \dots & m(p + q) - 2 & m(p + q) - 1 & m(p + q) \end{array}$$

We have $e_{mp,p-2} + e_{mp,p-1} + e_{mp,p} \leq (3m-1)p + 3mq - 1 \leq 3mp + 3mq - 3 = e_{mp+1,p-2} + e_{mp+1,p-1} + e_{mp+1,p}$. Since $e_{mp,g} < e_{mp+1,g}$ (in case there is no $e_{mp,g}$, we assume $e_{mp,g} = 0$), $1 \leq g \leq p-2$ and $p \geq 2$, therefore $wt(r_{mp}) < wt(T_m^t)$.

When the array L_j , $1 \leq j \leq m$, is removed from the construction given in the proof of Case 1 of Theorem 2.1, the sausage graph degenerates into a path and so gives an alternative proof of the path being antimagic. The path has been proved to be antimagic originally in [5, 8].

Corollary 2.2. *The generalized sausage graph $S(K_1, m) = P_{m+2}$, $m \geq 1$, is antimagic.*

Corollary 2.3. *The generalized sausage graph $S(nK_1, m)$, $m \geq 1$ and $n \geq 2$, is antimagic.*

PROOF. For $n = 2$, $S(2K_1, m)$ is a circle C_{2m+2} . It has been proved to be antimagic in [5, 8].

We next prove it for $n \geq 3$. We first label the edges of the path P_{m+2} as shown in the diagram in Figure 3. We label that the edge e_i , $1 \leq i \leq m+1$, of P_{m+2} labels with i . This ensures that the weights of the vertices with degree 2 are pairwise distinct. To build the graph $S(nK_1, m)$ we use n copies of P_{m+2} . Let L_j , $1 \leq j \leq n$, be the array of j -th copy of P_{m+2} , where the weights of the vertices with degree 2 are in the ascending order. We construct the array A of edge labels of $S(nK_1, m)$, $m \geq 1$ and $n \geq 2$, as follows.

- (1) Replace the label i of the edge e_i in the array L_j , $1 \leq j \leq n$, by adding $(j-1)(m+1)$ to each of the original edge labels;
- (2) Form the array A as shown below.

$$\begin{array}{c} L'_1 \\ L'_2 \\ \vdots \\ L'_n \\ A_1 \\ A_2 \end{array}$$

where L'_j , $1 \leq j \leq n$, is the array of the edge labels of the vertices of degree 2 of j th-copy of P_{m+2} , $A_1 = (m \ m + (m+1) \ m + 2(m+1) \ \dots \ m + (n-1)(m+1))$ and $A_2 = (m+1 \ 2(m+1) \ 3(m+1) \ \dots \ n(m+1))$.

We skip details of the proof when $n \geq 3$ and $m = 1$. For the case $n \geq 3$ and $m = 2$, we only need a small change from the case of $n = 3$ and $m = 2$ by swapping the labels 1 and 2; and the rest of the proof is skipped here since it is similar to the following case.

We now consider the case $n \geq 3$ and $m \geq 3$. By the construction of the array A , it clear that the weight of each vertex (row) in the array is less than the weight




FIGURE 3. Illustration of a labeling of the path P_{m+2} , $m \geq 1$

of the vertex (row) below, except the weight of the last four rows that need to be verified.

Let r_{nm-1} and r_{nm} be the last two rows of the subarray L'_n . We have $wt(r_{nm-1}) = m + (n - 1)(m + 1) + (m - 2) + (n - 1)(m + 1) = 2(n + 1)m - 4$, $wt(r_{nm}) = n(m + 1) + (m - 1) + (n - 1)(m + 1) = 2n(m + 1) - 2$, $wt(A_1) = nm + \frac{n(n-1)(m+1)}{2}$ and $wt(A_2) = \frac{n(n+1)(m+1)}{2}$.

For $n = 3$, we have $wt(r_{nm-1}) = 6m + 2$, $wt(r_{nm}) = 6m + 4$, $wt(A_1) = 6m + 3$ and $wt(A_2) = 6m + 6$.

For $n \geq 4$, we have $wt(A_1) - wt(r_{nm}) = \frac{n^2m - 3nm + n^2 - 5n + 4}{2} > 0$. Therefore, $wt(r_{nm-1}) < wt(r_{nm}) < wt(A_1) < wt(A_2)$.

We extend Theorem 2.1 to more general cases in the following theorems and corollaries.

Theorem 2.4. *Let $G \neq nK_1$, $n \geq 1$, be any connected or disconnected k -regular graph. Then the mixed generalized sausage graph $MS(G, m)$, $m \geq 3$, is antimagic.*

PROOF. Let L_j , $1 \leq j \leq m$, be the array of edge labels of the j -th copy of the graph G in $MS(G, m)$, $m \geq 3$. We construct the array A of edge labels of $MS(G, m)$, $m \geq 3$, as follows.

Case 1: $m \geq 3$ and m odd

Let T_l , $1 \leq l \leq 2m$, be the $(p \times 1)$ -array of the edges e_i , $1 \leq i \leq p$, where e_i are the edges of $MS(G, m)$, $m \geq 3$, that do not belong to any copy of G .

Subcase 1.1: $m = 3$

First consider K_p and then other graphs.

Subcase 1.1.1: $G = K_p$, $p \geq 2$

- (1) Label the edges e_i , $1 \leq i \leq p$, in the row i of the array T_l , $1 \leq l \leq 2m$, with $i + (l - 1)p$;
- (2) Replace the edge labels in the array L_j , $1 \leq j \leq m$, with new labels obtained by adding $2mp + (j - 1)q$ to each of the original edge labels;
- (3) Form the array A as shown below.

$$\begin{array}{cccc}
 & & T_1^t & T_3^t \\
 & & T_2^t & T_4^t \\
 & T_3 & T_5 & L_1 \\
 & T_4 & T_6 & L_2 \\
 T_1 & T_2 & T_5 & T_6 & L_3
 \end{array}$$

By the construction of the array A , it is clear that the weight of each vertex (row) in the array is less than the weight of the vertex (row) below, except possibly for some special cases below.

Let $wt(r_f)$ be the weight of the row r_f .

(a) $T_2^t T_4^t$ and the first row of $T_3 T_5 L_1$

Since the least possible edge labels (that yield the least possible weight) of the vertex in the array L_1 are $1+6p, 2+6p, \dots, k+6p$, hence $wt(r_3) > (1+2p) + (1+4p) + (1+6p) + (2+6p) + \dots + (k+6p) = 6p^2 + \frac{p^2-p}{2} + 2 > 5p^2 + p = wt(T_2^t T_4^t)$.

(b) Last row of $T_4 T_6 L_2$ and the first row of $T_1 T_2 T_5 T_6 L_3$

Let r_{2m+2} and r_{2m+3} be the last row of the array $T_4 T_6 L_2$ and the first row of the array $T_1 T_2 T_5 T_6 L_3$, respectively. We have the edge labels of the rows r_{2m+2} and r_{2m+3} as shown below.

$$\begin{array}{rcccc} r_{2m+2} : & & 4p & 6p & \dots \\ r_{2m+3} : & 1 & 1+p & 1+4p & 1+5p & \dots \end{array}$$

Since $4p+6p = 10p < 10p+4 = 1 + (1+p) + (1+4p) + (1+5p)$ and all edge labels in the array L_2 are less than the least possible edge label in the array L_3 , hence $wt(r_{2m+2}) < wt(r_{2m+3})$.

Subcase 1.1.2: $G \neq K_p$, $p \geq 1$

- (1) Replace the edge labels in the array L_j , $1 \leq j \leq m$, with new labels obtained by adding $(j-1)q$ to each of the original edge labels;
- (2) Label the edges e_i , $1 \leq i \leq p$, in the row i of the array T_l , $1 \leq l \leq 2m$, with $i + (l-1)p + mq$;
- (3) Form the array A as shown below.

$$\begin{array}{cccccc} & & L_1 & T_1 & T_3 & \\ & & & L_2 & T_2 & T_4 & \\ & & & & & & L_3 & T_3 & T_4 & T_5 & T_6 & \\ & & & & & & & & & T_1^t & T_5^t & \\ & & & & & & & & & T_2^t & T_6^t & \end{array}$$

By the construction of the array A , it is clear that the weight of each vertex (row) in the array is less the weight of the vertex (row) below, except the weights of the last row of the subarray $L_3 T_3 T_4 T_5 T_6$ and the row $T_1^t T_5^t$ that need to be verified.

Let r_{3m} and r_{3m+1} be the last row of the subarray $L_3 T_3 T_4 T_5 T_6$ and the row $T_1^t T_5^t$, respectively. Since $p \geq k+2$, we have the edge labels of the rows r_{3m} and r_{3m+1} as shown below.

$$\begin{array}{rcccccc} r_{3m} : & \dots & 3p+3q & 4p+3q & 5p+3q & 6p+3q \\ r_{3m+1} : & \dots & 5p+3q-3 & 5p+3q-2 & 5p+3q-1 & 5p+3q \end{array}$$

We have $(3p+3q) + (4p+3q) + (5p+3q) + (6p+3q) = 18p+12q < 20p+12q-6 = (5p+3q-3) + (5p+3q-2) + (5p+3q-1) + (5p+3q)$, for $p > 3$, and the largest possible of the rest of edge labels in the row r_{3m} is less than the least edge label of




FIGURE 4. Illustration of antimagic labeling of $MS(G, m)$, $m > 4$ and m odd

the rest of the edge labels in the row r_{3m+1} , then $wt(r_{3m}) < wt(r_{3m+1})$.

Subcase 1.2: $m > 4$

- (1) Replace the edge labels in the array L_j , $1 \leq j \leq m$, with new labels obtained by adding $(j - 1)(p + q)$ to each of the original edge labels;
- (2) Label the edges e_i , $1 \leq i \leq p$, in the row i of the array T_l , $1 \leq l \leq 2m$, with $i + (l - 1)p + lq$, for $1 \leq l \leq m$, and $i + (l - 1)p + mq$, for $m + 1 \leq l \leq 2m$;
- (3) Form the array A as shown below.

	L_1	T_1	T_3		
	L_2	T_2	T_4		
	L_3	T_3	T_5	T_{m+2}	
	L_4	T_4	T_6	T_{m+3}	
	\vdots	\vdots	\vdots	\vdots	
	L_{m-1}	T_{m-1}	T_{m+1}	T_{2m-2}	
	L_m	T_m	T_{m+1}	T_{2m-1}	T_{2m}
T_1^t	T_{m+2}^t	\cdots	\cdots	T_{2m-3}^t	T_{2m-1}^t
T_2^t	T_{m+3}^t	\cdots	\cdots	T_{2m-2}^t	T_{2m}^t

The diagram in Figure 4 illustrates the antimagic labeling used here.

By the construction of the array, it is clear that the weight of each vertex (row) in the array is less than the weight of the vertex (row) below, except the weights of the last row of the subarray $L_m T_m T_{m+1} T_{2m-1} T_{2m}$ and the row $T_1^t T_{m+2}^t \cdots T_{2m-5}^t T_{2m-3}^t T_{2m-1}^t$ that need to be verified.

Let r_{mp} and r_{mp+1} be the last row of the subarray $L_m T_m T_{m+1} T_{2m-1} T_{2m}$ and the row $T_1^t T_{m+2}^t \cdots T_{2m-5}^t T_{2m-3}^t T_{2m-1}^t$, respectively. We have the edge labels of the rows r_{mp} and r_{mp+1} as shown below.

$$\begin{array}{l}
 r_{mp} : \quad \cdots \quad (m + 1)p + mq \quad (2m - 1)p + mq \quad 2mp + mq \\
 r_{mp+1} : \quad \cdots \quad (2m - 1)p + mq - 2 \quad (2m - 1)p + mq - 1 \quad (2m - 1)p + mq
 \end{array}$$

We have $(m + 1)p + mq + (2m - 1)p + mq + 2mp + mq = 5mp + 3mq < (6m - 3)p + 3mq - 3 = (2m - 1)p + mq - 2 + (2m - 1)p + mq - 1 + (2m - 1)p + mq$

and the rest of the edge labels in the row r_{mp} is less than the edge label in the row r_{3m+1} of the corresponding column. Hence $wt(r_{mp}) < wt(r_{mp+1})$.

Case 2: $m \geq 4$ and m even

Let T_l , $1 \leq l \leq 2m - 1$, be the $(p \times 1)$ -array of the edges e_i , $1 \leq i \leq p$, where e_i are the edges of $MS(G, m)$, $m \geq 3$, that do not belong to any copy of G .

- (1) Replace the edge labels in the array L_j , $1 \leq j \leq m$, with new labels by adding $(j - 1)q$ to each of the original edge labels;
- (2) Label the edges e_i , $1 \leq i \leq p$, in the row i of the array T_l , $1 \leq l \leq 2m - 1$, with $i + (l - 1)p + mq$;
- (3) Form the array A into two cases as shown below.

Subcase 2.1: $m = 4$

$$\begin{array}{cccc} L_1 & T_1 & T_3 & \\ L_2 & T_2 & T_4 & \\ L_3 & T_3 & T_5 & T_6 \\ L_4 & T_4 & T_6 & T_7 \\ & & T_1^t & T_5^t \\ & & T_2^t & T_7^t \end{array}$$

For $G = K_2$, we swap the labels 17 and 18 in the array T_7 .

For $q \geq p$, by the construction of the array A , it is clear that the weight of each vertex (row) is less than the weight of the vertex (row) below, except the last row of the subarray $L_4T_4T_6T_7$ and the row $T_1^tT_5^t$ that need to be verified.

Let $e_{f,g}$ be the edge label in the row f and the column g . Let r_{4p} and r_{4p+1} be the last row of subarray $L_4T_4T_6T_7$ and the row $T_1^tT_5^t$, respectively. We have the edge labels in the rows r_{4p} and r_{4p+1} as shown below.

$$\begin{array}{cccc} r_{4p} : & & \dots & 6p + 4q & 7p + 4q \\ r_{4p+1} : & 1 + 4q & \dots & 5p + 4q - 1 & 5p + 4q \end{array}$$

Since $e_{4p,2p-1} + e_{4p,2p} = (6p + 4q) + (7p + 4q) = 13p + 8q < 10p + 12q = (1+4q) + (5p+4q-1) + (5p+4q) = e_{4p+1,1} + e_{4p+1,2p-1} + e_{4p+1,2p}$ and $e_{4p,g} < e_{4p+1,g}$, for $2p - (k + 2) \leq g \leq 2p - 2$, hence $wt(r_{4p}) < wt(r_{4p+1})$.




FIGURE 5. Illustration of antimagic labeling of $MS(G, m)$, $m > 4$ and m even

Subcase 2.2: $m \geq 6$

		L_1	T_1	T_3	
		L_2	T_2	T_4	
		L_3	T_3	T_5	T_{m+2}
		L_4	T_4	T_6	T_{m+3}
		\vdots	\vdots	\vdots	\vdots
		L_{m-1}	T_{m-1}	T_{m+1}	T_{2m-2}
		L_m	T_m	T_{2m-2}	T_{2m-1}
T_1^t	T_{m+1}^t	\dots	\dots	T_{2m-6}^t	T_{2m-4}^t
T_2^t	T_{m+3}^t	\dots	\dots	T_{2m-3}^t	T_{2m-1}^t

The diagram in Figure 5 illustrates the antimagic labeling used here.

By the construction of the array A , it is clear that the weight of each vertex (row) is less than the weight of the vertex (row) below, except the last row of the subarray $L_m T_m T_{2m-2} T_{2m-1}$ and the row $T_1^t T_{m+1}^t \dots T_{2m-8}^t T_{2m-6}^t T_{2m-4}^t$ that need to be verified.

Let $e_{f,g}$ be the edge label in the row f and the column g . Let r_{mp} and r_{mp+1} be the last row of subarray $L_m T_m T_{2m-2} T_{2m-1}$ and the row $T_1^t T_{m+1}^t \dots T_{2m-8}^t T_{2m-6}^t T_{2m-4}^t$, respectively. We have the greatest possible labels of the row r_{mp} (when the last row of L_m has the largest labels) and the labels of the r_{mp+1} as shown below. We consider in three cases.

(a) $m = 6$

$$\begin{array}{rcccccc}
 r_{mp} : & & \dots & mq & m(p+q) & (2m-2)p+mq & (2m-1)p+mq \\
 r_{mp+1} : & 1+mq & \dots & a-3 & a-2 & a-1 & a
 \end{array}$$

where $a = (2m-4)p + mq$.

Since, for $p \geq 2$, we have $\sum_{h=0}^3 e_{mp, \frac{m}{2}p-h} = mq + m(p+q) + ((2m-2)p + mq) + ((2m-1)p + mq) = (5m-3)p + 4mq < 4(2m-4)p + 4mq - 6 = a + (a-1) + (a-2) + (a-3) = \sum_{h=0}^3 e_{mp+1, \frac{m}{2}p-h}$; and $e_{mp,g} < e_{mp+1,g}$, for $\frac{m}{2}p - (k+2) \leq g \leq \frac{m}{2}p - 4$, hence $wt(r_{mp}) < wt(r_{mp+1})$.

Corollary 2.5. *The mixed generalized sausage graph $MS(K_1, m)$, $m \geq 3$, is antimagic.*

PROOF. We divide the proof into three cases.

Case 1: $m = 3$

The same construction as the one given in the proof of Subcase 1.1.1 of Theorem 2.4 also works whenever the array L_j , $1 \leq j \leq 3$, is removed.

Case 2: $m \geq 4$

Subcase 2.1: m is odd

The same construction as the one given in the proof of Subcase 1.2 of Theorem 2.4 also works whenever the array L_j , $1 \leq j \leq m$, is removed.

Subcase 2.2: m is even

The same construction as the one given in the proof of Case 2 of Theorem 2.4 also works whenever the array L_j , $1 \leq j \leq m$, is removed. Using this construction, when $m = 4$ and $m = 6$, there are some weights are equal. However, we need only a small change by swapping the labels 5 and 6, then all vertex weights are pairwise distinct.

Corollary 2.6. *The mixed generalized sausage graph $MS(nK_1, m)$, $m \geq 3$ and $n \geq 2$, is antimagic.*

PROOF. For m odd, the same constructions as the one given in the proof of Subcase 1.1.2 and Subcase 1.2 of Theorem 2.4 also work whenever the array L_j , $1 \leq j \leq m$, is removed.

For $m = 4$, the same construction as the one given in the proof of Subcase 2.1 of Theorem 2.4 also works whenever the array L_j , $1 \leq j \leq m$, is removed; except when $n = 2$, we need a small change by swapping 13 and 14, and when $n = 3$ by swapping 20 and 21.

For $m \geq 6$ and m even, the same construction as the one given in the proof of Case 2.2 of Theorem 2.4 also works whenever the array L_j , $1 \leq j \leq m$, is removed.

Recall the definition of the reverse T^\uparrow from Section 2; we will use it in the proofs of the following theorems and corollaries.

Theorem 2.7. *Let $G \neq nK_1$, $n \geq 1$, be any connected or disconnected k -regular graph. Then the complete mixed generalized sausage graph $CMS(G, m)$, $m \geq 3$, is antimagic.*

PROOF. Let L_j , $1 \leq j \leq m$, be the array of edge labels of the j -th copy of the graph G in $CMS(G, m)$, $m \geq 3$. Let T_l , $1 \leq l \leq 3m$, be the $(p \times 1)$ -array of the edges e_i , $1 \leq i \leq p$, where e_i are the edges of $CMS(G, m)$, $m \geq 3$, that do not belong to any copy of G . We construct the array A of $CMS(G, m)$, $m \geq 3$, as follows.




FIGURE 7. Illustration of antimagic labeling of $CMS(G, m)$, $m \geq 3$ and m even

- (1) Replace the edge labels in the array L_j , $1 \leq j \leq m$, with new labels obtained by adding $(j - 1)(p + q)$ to each of the original edge labels;
- (2) Label the edges e_i , $1 \leq i \leq p$, in the row i of the array T_l , $1 \leq l \leq 3m$, with $i + (l - 1)p + lq$, for $1 \leq l \leq m$, and $i + (l - 1)p + mq$, for $m + 1 \leq l \leq 3m$;
- (3) Form the array A as shown below.

For $m = 3, 4$,

L_1	T_1	T_3	T_5	T_6	L_1	T_1	T_3	T_5	T_6
L_2	T_2	T_4	T_5	T_7^\uparrow	L_2	T_2	T_4	T_6	T_8^\uparrow
L_3	T_3	T_4	T_8	T_9^\uparrow	L_3	T_3	T_5	T_9	T_{10}^\uparrow
		T_1^t	T_7^t	T_8^t	L_4	T_4	T_5	T_{11}	T_{12}^\uparrow
		T_2^t	T_6^t	T_9^t		T_1^t	T_8^t	T_9^t	T_{11}^t
						T_2^t	T_7^t	T_{11}^t	T_{12}^t

More generally, for $m > 4$,

			L_1	T_1	T_3	T_{m+2}	T_{m+3}
			L_2	T_2	T_4	T_{m+2}	T_{m+4}^\uparrow
			L_3	T_3	T_5	T_{m+5}	T_{m+6}
			L_4	T_4	T_6	T_{m+7}	T_{m+8}
			\vdots	\vdots	\vdots	\vdots	\vdots
			L_{m-2}	T_{m-2}	T_m	T_{3m-5}	T_{3m-4}
			L_{m-1}	T_{m-1}	T_{m+1}	T_{3m-3}	T_{3m-2}
			L_m	T_m	T_{m+1}	T_{3m-1}	T_{3m}^\uparrow
T_1^t	T_{m+4}^t	T_{m+5}^t	\dots	\dots	T_{3m-5}^t	T_{3m-3}^t	T_{3m-1}^t
T_2^t	T_{m+3}^t	T_{m+6}^t	\dots	\dots	T_{3m-4}^t	T_{3m-2}^t	T_{3m}^t

The diagram in Figure 7 illustrates the antimagic labeling used here.

By the construction of the array A , it clear that the weight of each vertex (row) is less than the weight of the vertex (row) below, except some special cases that need to be verified.

We skip details for the case $m = 3$ and next verify for the case $m \geq 4$.

Let $e_{f,g}$ be the label at the row f and the column g in the array A .

(a) Rows r_p and r_{p+1}

We have the edge labels of rows r_p and r_{p+1} as shown.

$$\begin{array}{l} r_p : \quad \dots \quad (m+2)p + mq \quad (m+3)p + mq \\ r_{p+1} : \quad \dots \quad 1 + (m+1)p + mq \quad (m+4)p + mq \end{array}$$

Since $e_{p,mp-1} + e_{p,mp} = (2m+5)p + 2mq < (2m+5)p + 2mq + 1$, for all p and q , and $e_{p,g} < e_{p+1,g}$, for $1 \leq g \leq mp-2$, hence $wt(r_p) < wt(r_{p+1})$.

(b) Rows $r_{(m-1)p}$ and $r_{(m-1)p+1}$

We have $e_{(m-1)p,mp-2} + e_{(m-1)p,mp} = (4m-1)p + 2mq < 4mp + 2mq + 1 = e_{(m-1)p+1,mp-2} + e_{(m-1)p+1,mp}$, for all p and q ; and $e_{(m-1)p,g} < e_{(m-1)p+1,g}$, for $1 \leq g \leq mp-3$, and $g = mp-1$. Then $wt(r_{(m-1)p}) < wt(r_{(m-1)p+1})$.

(c) Rows r_{mp} and r_{mp+1}

Since $m \geq 4$ and $p \geq 2$, it is clear that $r_{mp} < r_{mp+1}$.

(d) Rows r_{mp+1} and r_{mp+2}

Let A and B be the sum of all the edge labels in subarrays $T_1^t T_{m+4}^t$ and $T_2^t T_{m+3}^t$, respectively. It is easy to check that $A < B$ and $e_{mp+1,g} < e_{mp+2,g}$, for $2p+1 \leq g \leq mp$. Hence $wt(r_{mp+1}) < wt(r_{mp+2})$.

The same construction as the one given in the proof of Theorem 2.7 also works when the array L_j , $1 \leq j \leq m$, is removed. We have

Corollary 2.8. *The complete mixed generalized sausage graph $CMS(nK_1, m)$, $m \geq 3$, $n \geq 1$, is antimagic.*

See Section 2 for definition of $CMS^-(G, m)$, then we have

Corollary 2.9. *Let $G \neq nK_1$, $n \geq 1$, be any connected or disconnected k -regular graph. Then the complete mixed generalized sausage graph $CMS^-(G, m)$, $m \geq 2$, is antimagic.*

PROOF. If $m \geq 3$, remove T_{m+2} from the array A in the proof of Theorem 2.7 and replace the array T_l with T_{l-1} , for $m+3 \leq l \leq 3m$.

If $m = 2$, we construct the array A as shown.

$$\begin{array}{cccc} L_1 & T_1 & T_2 & T_3 \\ L_2 & T_1 & T_4 & T_5^\uparrow \\ & & T_2^t & T_4^t \\ & & T_3^t & T_5^t \end{array}$$

For $p = 2$, that is, $G = K_2$, it is easy to check that $CMS^-(K_2, 2)$ is antimagic.

For $p \geq 3$, it is similar to (a) in the proof of Theorem 2.7 for checking $wt(r_p) < wt(r_{p+1})$. We next prove that $wt(r_{2p}) < wt(r_{2p+1})$. We have the edge labels of r_{2p} and $wt(r_{2p+1})$ as shown.

$$\begin{array}{rcccc} r_{2p} : & \dots & p+q & 4p+3q & 1+4p+3q \\ r_{2p+1} : & \dots & 4p+3q-2 & 4p+3q-1 & 4p+3q \end{array}$$

Since $e_{2p,2p-2} + e_{2p,2p-1} + e_{2p,2p} = 9p + 7q + 1 < 12p + 9q - 3 = e_{2p+1,2p-2} + e_{2p+1,2p-1} + e_{2p+1,2p}$ and $e_{2p,g} < e_{2p+1,g}$, for $2p - k - 2 \leq g \leq 2p - 3$, hence $wt(r_{2p}) < wt(r_{2p+1})$.

The same construction as the one given in Corollary 2.9 also works when the array L_j , $1 \leq j \leq m$, is removed, except when $m = 2$ and $n = 1$, it needs a small change by swapping the labels 1 and 2. Then we have

Corollary 2.10. *The complete mixed generalized sausage graph $CMS^-(nK_1, m)$, $m \geq 2$, $n \geq 1$, is antimagic.*

REFERENCES

- [1] Alon, N., Kaplan, G., Lev, A., Roditty, Y. and Yuster, R., “Dense graphs are antimagic”, *J. Graph Theory*, **47**(4) (2004), 297–309.
- [2] Bača, M. and Miller, M., *Super Edge-Antimagic Graphs: A Wealth of Problems and Some Solutions*, BrownWalker Press, Boca Raton, Florida, USA, 2008.
- [3] Bača, M., Lin, Y. and Semaničová-Feňovčíková, A., “Note on super antimagicness of disconnected graphs”, *Int. J. Graphs and Comb.*, **6**(1) (2009), 47–55.
- [4] Gallian, J. A., “A Dynamic survey of graph labeling”, *Electron. J. Combin.*, **19**(#DS6) (2012).
- [5] Hartsfield, N. and Ringel, G., *Pearls in graph theory: A comprehensive introduction*, Academic Press Inc., Boston, MA, 1990.
- [6] Miller, M., Phanalasy, O., Ryan, J. and Rylands, L., “Antimagicness of some families of generalized graphs”, *Austral. J. Combin.* **53** (2012), 179–190.
- [7] Phanalasy, O., Miller, M., Rylands, L. and Lieby, P., “On a relationship between completely separating systems and antimagic labeling of regular graphs”, *LNCS* **6460** (2011), 238–241.
- [8] Ryan, J., Phanalasy, O., Miller, M. and Rylands, L., “On antimagic labeling for generalized web and flower graphs”, *LNCS* **6460** (2011), 303–313.
- [9] Rylands, L., Phanalasy, O., Ryan, J. and Miller, M., “An application of completely separating systems to graph labeling”, *LNCS* **8288** (2013), 376–387.
- [10] Zhang, Y. and Sun, X., “The antimagicness of the Cartesian product of graphs”, *Theoretical Computer Science*, **410** (2009), 727–735.