

CORRIGENDUM TO NEW INEQUALITIES ON
HOMOGENEOUS FUNCTIONS, J. INDONES. MATH.
SOC. 15 (2009), NO. 1, 49-59

V. LOKESHA¹, K.M. NAGARAJA², AND Y. SIMSEK³

¹ Department of Mathematics, Acharya Institute of Technology,
Soldevana- halli, Bangolore-90, India
lokiv@yahoo.com

²Department of Mathematics, Sri Krishna Institute of Technology,
Chikkabanavara, Hesaraghata Main Road, Karnataka, Bangolore-90, India
2406@yahoo.co.in

³ University of Akdeniz, Faculty of Arts and Science,
Department of Mathematics, 07058, Antalya, Turkey
ysimsek@akdeniz.edu.tr

The paper contains typing errors.

Theorem 2 Let $\mu_1, \mu_2 \in (-2, \infty)$, $r < 1$. If $\mu_1 \leq \frac{4}{1-r} \leq \mu_2$, then

$$gn_{\mu_2, r}(a, b) \leq L(a, b) \leq Gn_{\mu_1, r}(a, b). \quad (12)$$

Furthermore $\mu_1 = \mu_2 = \frac{4}{1-r}$ is the best possibility for inequality (12). Also for $r = 0$,

$$gn_{\mu_2, 0}(a, b) \leq L(a, b) \leq Gn_{\mu_1, 0}(a, b). \quad (13)$$

Furthermore $\mu_1 = \mu_2 = 4$ is the best possibility for inequality (13).

Theorem 3 For $\mu_1, \mu_2 \in (-2, \infty)$, $r \neq \frac{2}{3}$, $r < 1$ and if $\mu_1 \leq \frac{2}{2-3r} \leq \mu_2$, then

$$gn_{\mu_2, r}(a, b) \leq I(a, b) \leq Gn_{\mu_1, r}(a, b). \quad (15)$$

Furthermore $\mu_1 = \mu_2 = \frac{2}{2-3r}$ is the best possibility for inequality (15). Also for $r = 0$,

$$gn_{\mu_2, 0}(a, b) \leq I(a, b) \leq Gn_{\mu_1, 0}(a, b). \quad (16)$$

Furthermore $\mu_1 = \mu_2 = 1$ is the best possibility for inequality (16).

Theorem 4 For $\mu_1, \mu_2 \in (-2, \infty)$, $r \neq 0$ and if $\mu_2 \leq \frac{2}{r} - 2 \leq \mu_1$, then

$$gn_{\mu_2, 0}(a, b) \leq M_r(a, b) \leq Gn_{\mu_1, 0}(a, b). \quad (17)$$

Furthermore $\mu_1 = \mu_2 = \frac{2}{r} - 2$ is the best possibility for inequality (17)

The above Theorems should be corrected to as follows:

Theorem 2 For $r \neq \frac{1}{3}$ and $\mu_1, \mu_2 \in (-2, \infty)$ such that $\mu_1 \leq \frac{4}{1-3r} \leq \mu_2$, then

$$gn_{\mu_2, r}(a, b) \leq L(a, b) \leq Gn_{\mu_1, r}(a, b). \quad (12)$$

Furthermore $\mu_1 = \mu_2 = \frac{4}{1-3r}$ is the best possibility for inequality (12). Also for $r = 0$,

$$gn_{\mu_2, 0}(a, b) \leq L(a, b) \leq Gn_{\mu_1, 0}(a, b). \quad (13)$$

Furthermore $\mu_1 = \mu_2 = 4$ is the best possibility for inequality (13).

Theorem 3 For $r \neq \frac{2}{3}$ and $\mu_1, \mu_2 \in (-2, \infty)$ such that $\mu_1 \leq \frac{2}{2-3r} \leq \mu_2$, then

$$gn_{\mu_2, r}(a, b) \leq I(a, b) \leq Gn_{\mu_1, r}(a, b). \quad (15)$$

Furthermore $\mu_1 = \mu_2 = \frac{2}{2-3r}$ is the best possibility for inequality (15). Also for $r = 0$,

$$gn_{\mu_2, 0}(a, b) \leq I(a, b) \leq Gn_{\mu_1, 0}(a, b). \quad (16)$$

Furthermore $\mu_1 = \mu_2 = 1$ is the best possibility for inequality (16).

Theorem 4 For $r \neq 1$ and $\mu_1, \mu_2 \in (-2, \infty)$ such that $\mu_2 \leq \frac{r}{1-r} \leq \mu_1$, then

$$gn_{\mu_2, 0}(a, b) \leq M_r(a, b) \leq Gn_{\mu_1, 0}(a, b). \quad (17)$$

Furthermore $\mu_1 = \mu_2 = \frac{r}{1-r}$ is the best possibility for inequality (17).

Remark. Carlson [1] and Lin [2] gave some inequalities on mean and logarithmic mean.

REFERENCES

- [1] Carlson, B.C., "The logarithmic mean", *Amer. Math. Monthly*, **79** (1972), 615-618.
- [2] Lin, T.P., "Mean and logarithmic mean", *Amer. Math. Monthly*, **81** (1974), 879-883.